

# **Disease Control Priorities Project**

## **Working Paper No. 30** **March 2003**

### **Background Paper on the Neurobiology of Nicotine Addiction**

**Athina Markou<sup>1</sup>**

**George F. Koob<sup>1</sup>**

**Jack E. Henningfield<sup>2</sup>**

<sup>1</sup> Department of Neuropharmacology, The Scripps Research Institute, La Jolla, California, U.S.A.

<sup>2</sup> Department of Psychiatry and Behavioral Sciences, Johns Hopkins University School of Medicine, Baltimore, Maryland, U.S.A.; Research and Health Policy, Pinney Associates, Bethesda, Maryland, U.S.A.

Correspondence: Dr. Athina Markou  
Department of Neuropharmacology, CVN-7  
The Scripps Research Institute  
10550 North Torrey Pines Road  
La Jolla, CA 92037  
tel: 858-784-7244  
fax: 858-784-7405  
email: amarkou@scripps.edu

The Disease Control Priorities Project is a joint effort of The World Bank, the Fogarty International Center of the National Institutes of Health, the Bill & Melinda Gates Foundation, and the World Health Organization.



*Disease Control Priorities Project*  
*Nicotine Addiction Workshop: Mumbai, India, 10-11 March 2003*

**Background paper on the neurobiology of nicotine addiction**

Since the 1980s, it has become increasingly recognized by clinicians, researchers, and public health experts that tobacco products are appropriately categorized among the most addictive and deadly of all dependence producing substances (World Health Organization 2001; Royal College of Physicians of London 2000). There are over 4000 chemicals in cigarette smoke, many of which could potentially contribute to the addictive properties of tobacco. In light of myriad preclinical studies demonstrating nicotine's reinforcing properties in many species including humans (Goldberg et al. 1981; Risner & Goldberg 1983; Goldberg & Henningfield 1988; Corrigall & Coen 1991; Fudala et al. 1985; Huston-Lyons & Kornetsky 1992; Watkins et al. 1999; Markou & Paterson 2001), there is little debate about the fact that people smoke primarily to experience the acute psychopharmacological properties of nicotine, and that nicotine is one of the major components in tobacco smoke responsible for addiction (Balfour 1984; USDHHS 1988; Royal Society of Canada 1989; Stolerman 1991; Stolerman & Jarvis 1995; Royal College of Physicians of London 2000). Nevertheless, in addition to nicotine, there are other factors that contribute to the tobacco smoking habit, as suggested by the fact that tobacco craving is only partially relieved by administration of pure nicotine forms. These other factors include sensory (e.g., smell and taste of cigarettes) and conditioned stimuli (e.g., the sight of other people smoking, tobacco advertisements), as well as other ingredients in tobacco smoke (e.g., tar) that may have synergistic effects with nicotine (Butschsky et al. 1995; Rose et al. 1993; Shahan et al. 1999; Crooks & Dwoskin 1997; Jacob et al. 1999; Bardo et al. 1999; Caggiula et al. 2001a, 2001b; Perkins et al. 2001). Thus, it is important to emphasize here that although pure nicotine can serve as a positive reinforcer, it does not mean that the reinforcing effects of cigarette smoke are fully accounted for by nicotine. As indicated above, sensory effects of tobacco can acquire conditioned reinforcing properties. In addition, other chemicals present in smoke can interact with nicotine to enhance nicotine's reinforcing effects by either having their own primary pharmacological effects, in addition to potentially influencing nicotine dose delivered and absorbed, and the speed of nicotine delivery. For example, Philip Morris researchers found that acetaldehyde (which is present in cigarette smoke) can serve as a reinforcer in its own right for animals (DeNobel & Mele as reported by the U.S. Food and Drug Administration 1996; Royal

College of Physicians of London 2000). Further, these investigators found that when nicotine and acetaldehyde were administered simultaneously, the reinforcing effects were stronger than those produced by either drug alone. Other substances also have been investigated such as the tobacco alkaloid and active nicotine metabolite nornicotine which is self-administered in rats (Bardo et al. 1999), and which also produces psychomotor stimulant and discriminative stimulus effects (Bardo et al. 1997; Dwoskin et al. 1999). Such work has important implications for understanding the powerfully addicting effects of tobacco (Royal College of Physicians of London 2000). The present paper will focus on the neuropharmacology of nicotine, and on the neuronal substrates mediating the reinforcing effects of acute nicotine and nicotine withdrawal partly because in neurobiological research it is important to isolate and study each factor thoroughly before interactions of various factors are investigated, and because nicotine is probably one of the main ingredients in tobacco smoking leading to addiction. Similarly to other major drugs of abuse, it is believed that both the reinforcing properties of nicotine and the nicotine withdrawal syndrome lead to the development and maintenance of the tobacco smoking habit and the precipitation of relapse after a period of abstinence (Wikler 1973; Levine 1974; Stewart et al. 1984; Ludwig 1986; McLellan et al. 1986; Childress et al. 1986, 1988; Shiffman 1989; O'Brien et al. 1990; Koob et al. 1993; Markou et al. 1993, 1998; Hughes & Hatsukami 1992; Watkins et al. 2000a; Kenny & Markou 2001).

### **Nicotinic Acetylcholine Receptors**

Nicotine is an alkaloid that is present in concentrations of about 1-3% in tobacco cultivated for product manufacture (Browne 1990). Nicotine is an agonist at several subpopulations of nicotinic acetylcholine receptors (nAChR; Paterson & Nordberg 2000; Vidal 1996; Henningfield et al. 1996). Nicotinic receptors are expressed on mature skeletal muscle, in autonomic ganglia and within the central nervous system (Holladay et al. 1997). Not surprisingly, most interest in the behavioral actions of nicotine has focused on the role of nAChRs located within the central nervous system. Neuronal nAChRs, similar to other ligand-gated ion channels, are composed of five membrane spanning subunits that combine to form a functional receptor (Albuquerque et al. 1997; Dani 2000; Lindstrom et al. 1996; Lena & Changeux 1998; Role & Berg 1996). Individual neuronal nAChR subunits arrange in different combinations to form individual nAChRs with distinct pharmacological and kinetic properties. The neuronal  $\alpha$  subunit exists in nine isoforms ( $\alpha 2$ - $\alpha 10$ ) whereas the neuronal  $\beta$  subunit exists in three isoforms ( $\beta 2$ - $\beta 4$ ) (Elgoyhen et al. 2001; Arneric et al. 1995; Wonnacott 1997). Oocyte expression systems injected with pairwise combinations of different neuronal  $\alpha$  and  $\beta$  subunits have provided evidence that these subunits combine with a stoichiometry of  $2\alpha:3\beta$  to produce a functional neuronal nicotinic hetero-oligomeric receptor (Deneris et al. 1991; Conroy & Berg,

1995; Colquhoun & Patrick 1997). In contrast,  $\alpha 7$ ,  $\alpha 8$  and  $\alpha 9$  subunits form homo-oligomeric complexes composed of five  $\alpha$  subunits and lacking  $\beta$  subunits (Chen et al., 1998), with only the  $\alpha 7$  pentamer being expressed in the central nervous system.

Based on radioligand binding studies, neuronal nAChRs have been divided broadly into three classes within the rat brain: those with a high affinity binding site for [ $^3\text{H}$ ]-nicotine which correspond to  $\alpha 4$ -containing nAChRs, the  $\alpha 4\beta 2$  combination being the most abundant (Flores et al. 1992; Picciotto et al. 1995); those with high affinity for [ $^{125}\text{I}$ ]- $\alpha$ -bungarotoxin which correspond to  $\alpha 7$  nAChRs (Clarke 1992), and those with high affinity for neuronal bungarotoxin which correspond to  $\alpha 3$ -containing nAChRs (Schulz et al. 1991). The precise combinations of nAChR subunits that constitute active nAChRs within the CNS *in vivo* are unknown and have so far only been inferred by their pharmacological profile (Kaiser et al. 1998; Luo et al. 1998; Sershen et al. 1997; Sharples et al. 2000). However, with the advent of more sophisticated tools it is becoming possible to identify the nAChR subunits expressed by individual neurons within specific brain regions (Sheffield et al. 2000; Lena et al. 1999).

It has been proposed that the exclusive or predominant role of nAChRs in the brain is the modulation of neurotransmitter release (Wonnacott 1997). Brain nicotinic receptors are situated mainly on presynaptic terminals (Wonnacott 1997) but also are found at somatodendritic, axonal and postsynaptic locations (for review, Sargent 1993). Accordingly, by actions at nAChRs, nicotine stimulates the release of most neurotransmitters throughout the brain (Albuquerque et al. 1997; Alkondon et al. 1997; Gray et al. 1996; McGehee et al. 1995; McGehee & Role 1995; Role & Berg, 1996; Kenny et al. 2000; Araujo et al. 1988; Toide & Arima 1989; Wilkie et al. 1996; Grady et al. 2001). Therefore, as discussed below, it is likely that various neurotransmitter systems are involved in the adaptations that occur in response to chronic nicotine exposure that give rise to dependence and withdrawal responses.

### **Neurosubstrates of Nicotine Reinforcement**

Nicotine is an effective reinforcer for both humans and experimental animals as demonstrated by reports of mild euphoria in humans (Pomerleau & Pomerleau 1992) and intravenous self-administration studies in several species, such as rats, mice, nonhuman primates and humans (e.g., Goldberg et al. 1981, 1983; Henningfield et al. 1985; Goldberg & Henningfield, 1988; Corrigall & Coen 1989; Donny et al. 1995; Shoaib et al. 1997; Watkins et al. 1999; Picciotto et al. 1998; Markou & Paterson 2001). The mesolimbic dopaminergic system is considered of importance for the reinforcing properties of acute nicotine (for reviews, Watkins et al. 2000a; Picciotto & Corrigall 2002). Acute nicotine administration increased the firing rate of ventral tegmental area (VTA) dopaminergic neurons (Grenhoff et al. 1986; Pidoplichko et al. 1997), and elevated dialysate dopamine levels specifically in the shell of the nucleus accumbens

(Imperato et al. 1986; Damsma et al. 1989; Mifsud et al. 1989; Benwell & Balfour 1992; Pontieri et al. 1996; Nisell et al. 1997; Carboni et al. 2000), possibly through excitatory actions at nAChRs on mesolimbic dopaminergic neurons at both the VTA and the nucleus accumbens (Nisell et al. 1995; Teng et al. 1997; McGehee & Role 1996). Interestingly, nAChRs in the VTA were shown to play a more important role than those in the nucleus accumbens in mediating the effects of nicotine on dopamine release (Nisell et al. 1994a, b, 1997). The role of nAChRs on VTA dopaminergic neurons in nicotine reinforcement is supported by the finding that injections of the competitive nAChR antagonist dihydro- $\beta$ -erythroidine (DH $\beta$ E) (Williams & Robinson 1984) into the VTA (Corrigall et al. 1994), or lesions of the mesolimbic dopaminergic projections with 6-hydroxydopamine (Corrigall et al. 1992) or systemic administration of dopamine receptor antagonists (Corrigall & Coen 1991) decreased nicotine self-administration. Nevertheless, a dopamine receptor antagonist did not reverse nicotine-induced cognitive improvements demonstrating the specificity of the above results to the reinforcing effects of nicotine (Levin 1997; Levin & Simon 1998). Other mechanisms by which nicotine may elevate striatal dopamine levels are through enhancement of frontal cortex glutamatergic stimulation of ventral striatum dopamine release, and/or glutamatergic stimulation of VTA dopaminergic neurons projecting to the striatum. Nicotine increases glutamate release through agonism at excitatory presynaptic nAChRs on glutamatergic terminals (Toth et al. 1992; Vidal & Changeux 1993; Nisell et al. 1995; Radcliffe & Dani 1998; Damaj et al. 1999; Gioanni et al. 1999; Fu et al. 2000). Acute nicotine administration increased the release of glutamate in various brain sites including the VTA (Mansvelder and McGehee, 2000; Fu et al., 2000; Grillner and Svensson, 2000), nucleus accumbens (Reid et al. 2000), prefrontal cortex (Gioanni et al. 1999) and hippocampus (Gray et al. 1996). In terms of the VTA, it is believed that nicotine acts at presynaptic  $\alpha 7$  nAChRs located upon glutamate afferents (Mansvelder & McGehee 2000) to increase glutamate release in the VTA which in turn stimulates dopamine release in the nucleus accumbens (Nisell et al. 1994a; 1994b; Schilstrom et al. 1998a, b; Mansvelder & McGehee 2000; Fu et al. 2000). This enhanced glutamate release then acts at *N*-methyl-D-aspartate (NMDA) and non-NMDA receptor sites on postsynaptic dopamine neurons and increases their firing rate. Accordingly, NMDA receptor antagonists blocked tolerance to the locomotor depressant effects of acute nicotine (Shoaib & Stolerman 1992; Shoaib et al. 1994), and sensitization to the locomotor stimulant effects of chronic nicotine (Shoaib & Stolerman 1992). Blockade of postsynaptic metabotropic glutamate receptors 5 led to decreases in nicotine self-administration in rats and mice (Paterson et al. 2003), possibly by decreasing nicotine-stimulated dopamine release in the mesolimbic system. Similar decreases in nicotine self-administration are seen also after administration of dopaminergic and nAChR antagonists (Corrigall & Coen 1989, 1991; Watkins et al. 1999). In terms of nAChRs, several studies suggest an important involvement of the  $\alpha 4\beta 2$  and  $\alpha 7$  nAChR subtypes in both the

nicotine-induced release of dopamine and nicotine reinforcement (Picciotto et al. 1998; Schilstrom et al. 1998a; Watkins et al. 1999; Sharples et al. 2000; Grillner & Svensson 2000; Markou & Paterson 2001; however, see Grottick et al. 2000). Both the  $\alpha 4\beta 2$  and  $\alpha 7$  subtypes have been implicated also in nicotine's effects on memory (Bancroft & Levin, 2000; Levin et al. 1999), as well as the anxiolytic effects of nicotine (Cheeta et al. 2001; Gordon 1999) that also contribute to persistent tobacco use (USDHHS 1988).

In addition to dopamine and glutamate, another neurotransmitter system that may be critically involved in the reinforcing effects of acute nicotine is  $\gamma$ -aminobutyric acid (GABA). The dopaminergic neurons projecting from the VTA to the nucleus accumbens (Fallon & Moore 1978) receive descending GABAergic input from the ventral pallidum and the nucleus accumbens (Walaas & Fonnum 1979; Sugita et al. 1992). These GABAergic neurons have an inhibitory effect on dopaminergic tone (Klitenick et al. 1992; Engberg et al. 1993) at the level of both the VTA and the nucleus accumbens (Heimer et al. 1991; Heimer & Alheid 1991; Churchill et al. 1992; Kalivas 1993; Dewey et al. 1992; Kalivas et al. 1992). There are GABA inhibitory afferents to dopaminergic VTA neurons (e.g., Walaas & Fonnum 1980; Yim & Mogenson 1980), inhibitory GABA interneurons within the VTA, and medium spiny GABA neurons in the nucleus accumbens that also inhibit mesolimbic dopamine release (for review, Kalivas et al. 1992). Accordingly, enhancement of GABAergic transmission abolished both dopamine increases in the nucleus accumbens and the reinforcing effects of drugs of abuse, including those induced by nicotine (Dewey et al. 1999). Specifically, systemic injections of  $\gamma$ -vinyl GABA (GVG, also referred to as vigabatrin, an irreversible inhibitor of GABA transaminase, the primary enzyme involved in GABA metabolism; Jung et al. 1977; Lippert et al. 1977) that leads to increased GABA levels decreased nicotine self-administration in rats (Paterson & Markou 2002), and abolished the expression and acquisition of nicotine-induced conditioned place preference (Dewey et al. 1998). Further, GVG administration dose- and time-dependently lowered the nicotine-induced increases in nucleus accumbens dopamine in both naïve and chronically nicotine-treated rats as measured by *in vivo* microdialysis, and abolished nicotine-induced increases in dopamine in the striatum of primates as measured by positron emission tomography (Krystal et al. 2002). The use of receptor-selective agonists has suggested the involvement of GABA<sub>B</sub> receptors in these effects. Systemic or microinjections of baclofen or (3-amino-2[S]-hydroxypropyl)-methylphosphinic acid (CGP44532), two GABA<sub>B</sub> receptor agonists, systemically or directly into the nucleus accumbens shell, VTA or peduncular pontine nucleus, but not into the caudate-putamen, decreased the reinforcing effects of nicotine (Shoaib et al. 1998; Corrigall et al. 2000, 2001; Fattore et al. 2002; Paterson et al. 2003). Taken together, these results suggest that enhancement of GABA transmission through activation of GABA<sub>B</sub> receptors may block the reinforcing effects of nicotine. Interestingly, however, a human clinical study showed that a

single acute dose of baclofen had no effect on either the number of cigarettes smoked or craving for nicotine (Cousins et al. 2001). Nevertheless, other clinical studies using chronic administration of baclofen showed that baclofen reduced cocaine and alcohol abuse, and cue-induced brain activation (Ling et al. 1998; Addolorato et al. 2000, 2002a, b; Childress et al. 2002). Therefore, it is possible that chronic treatment with these GABAergic drugs is required before tobacco smoking is decreased.

Acute nicotine exposure has effects also on the hypothalamic-pituitary-adrenal (HPA) axis in rodents and humans that are hypothesized to be mediated by corticotropin-releasing factor (CRF) transmission in the paraventricular nucleus of the hypothalamus. In humans, two regular strength cigarettes (equivalent to 2 mg of nicotine) or intravenous nicotine increased corticosterone and adrenocorticotrophic hormone (ACTH) levels (Wilkins et al. 1982; Seyler et al. 1984; Stalke et al. 1992; Newhouse et al. 1990). Similarly, rats injected with nicotine (Cam et al. 1979; Andersson et al. 1983; Conte-Devolx et al. 1985) or exposed to inhalation of cigarette smoke showed increased levels of ACTH and corticosterone; this effect is mediated by central, but not peripheral nAChRs (Matta et al. 1987). Specifically, it was shown that the nAChR antagonist mecamylamine injected into the nucleus of the solitary tract prevented this brainstem nucleus from releasing norepinephrine into the paraventricular nucleus which in turn prevented HPA axis activation; this effect was not seen when mecamylamine was injected directly into the paraventricular nucleus (Matta et al. 1993). Several lines of evidence suggest that CRF in the paraventricular nucleus mediates the effects of nicotine on the HPA stress response system. Nicotine induces CRF release *in vitro* from both the rat hypothalamus (Hillhouse & Milton 1989) and a cell line derived from rat amygdala (Kasckow et al. 1999). Further, nAChRs are co-localized with CRF-positive vesicles on axon terminals in the paraventricular nucleus, demonstrating that nicotine could act directly on hypothalamic neurons to release CRF (Okuda et al. 1993). Finally, nicotine treatment induced c-fos protein expression in CRF-positive neurons in several brain areas, including the paraventricular nucleus (Valentine et al. 1996), bed nucleus of the stria terminalis, central nucleus of the amygdala, and the dorsal raphe (Matta et al. 1997), indicating that CRF may play a role in mediating the effect of nicotine on behaviors relating to stress and anxiety.

Finally, opioid receptor antagonists such as naloxone and naltrexone have been reported to modulate cigarette consumption and have been used as smoking cessation aids (Karras & Kane 1980; Wewers et al. 1998; Covey et al. 1999; but see Nemeth-Coslett & Griffiths 1986; Sutherland et al. 1995) suggesting that opioid receptors also may modulate the reinforcing effects of nicotine.

### **Adaptations in nAChR Function**



Chronic nicotine administration leads to an interesting paradoxical change in the function of nAChRs that consists of receptor desensitization leading to a receptor upregulation (Marks et al. 1983; Schwartz & Kellar 1983; Changeux et al. 1984; Wonnacott 1990; Flores et al. 1997; Perry et al. 1999; Mansvelder et al. 2002). There have been complex theoretical speculations about the role of nAChR desensitization and upregulation in the subjective effects of acute nicotine, and in the development and maintenance of nicotine dependence (Marks et al. 1992; Dani & Heinemann 1996; Dani et al. 2000; Quick & Lester 2002; Buisson & Bertrand 2002). Experimental evidence indicates that long-term exposure to nicotine induces an increase in the number (Buisson & Bertrand 2001; Marks et al. 1992; Wonnacott 1990) and function (Rowell & Wonnacott 1990) of nAChRs. Nevertheless, this finding is not consistent, and others see a decrease in nAChR number (Gentry et al. 2003) and function (Marks et al. 1993) with chronic exposure to nicotine. Further, behavioral findings are most readily explained by decreased number and/or function of nAChRs with the development of nicotine dependence (Epping-Jordan et al. 1998; Watkins et al. 2000b; Markou & Paterson 2001; Skjei & Markou 2003) that develop to counteract the continuous agonist actions of nicotine on the receptors. Most studies reporting changes in nAChR number and function have been conducted in *in vitro* experimental systems, and it is acknowledged that the functional significance of these changes *in vivo* are unknown (Buisson & Bertrand 2002) and need to be explored (Kellar et al. 1999). Further, it is known that different subtypes of nAChRs desensitize and upregulate at different rates which may explain the seemingly opposite effects seen in some cases (Kellar et al. 1999; Levin 2002; Buisson & Bertrand 2002). Thus, behaviorally we observe the net effect of these complex adaptations in different receptor types and brain sites. Adaptations in nAChR number and function may be long-lasting and may contribute to the difficulty many people have in achieving and sustaining nicotine abstinence even after the overt signs of nicotine withdrawal have subsided (see below). Although many smokers attempt to quit smoking and may be successful early on, only 20% of these individuals are still abstinent after one year (Hunt et al. 1971; Hunt & Besspalec 1974; Hughes et al. 1991a, b, 1992). These adaptations in nAChRs are likely to also mediate the rapid and high degree of tolerance seen to several of the effects of acute nicotine (Langley 1905; Royal College of Physicians of London 2000). Some degree of tolerance is gained also during each day of smoking and lost during the approximately 8 hours of tobacco deprivation during sleeping hours (Swedberg et al. 1990). The time course of gain and loss of tolerance varies across nicotine-induced responses (Collins & Marks 1989; Balfour & Fagerstrom, 1996; Royal College of Physicians of London 2000), with a large degree of tolerance to the subjective and cardiovascular effects of nicotine (Soria et al. 1996; USDHHS 1988; Royal College of Physicians of London 2000; Heishman & Henningfield 2000).

## **Neurosubstrates of Nicotine Withdrawal**

Smoking cessation leads to an aversive withdrawal syndrome in humans (Hughes et al. 1991b; Shiffman & Jarvik 1976), components of which are exhibited for 1-10 weeks post-smoking (Hughes 1992). This syndrome arises, at least in part, because of the cessation of administration of nicotine through tobacco smoking. The fact that cessation of nicotine administration leads to this syndrome is suggested by the fact that the withdrawal signs can be mitigated largely by administration of pure nicotine in a variety of forms (e.g., gum, patch, nasal delivery; Shiffman et al. 1998; Heishman et al. 1994; Pickworth et al. 1995; Hughes et al. 1990). Conversely, reduction of the nicotine content in smoked tobacco induced a withdrawal syndrome in smokers that was accompanied by a significant reduction in plasma nicotine levels (West et al. 1984). This nicotine withdrawal syndrome is comprised of "physical" or somatic, affective and cognitive components. The most common somatic symptoms include bradycardia and gastrointestinal discomfort. Affective symptoms primarily include depressed mood, dysphoria, craving, anxiety and irritability. A cognitive symptom of withdrawal is difficulty concentrating, while yet another symptom is increased appetite (American Psychiatric Association 1994; Hughes et al. 1991b; West et al. 1991; Glassman et al. 1990; Parrott 1993).

Several rodent models have been developed that serve to investigate the neurobiology of nicotine withdrawal and potentially evaluate medications for treating withdrawal. One of the first and most widely used measures of nicotine withdrawal in rodents is the measurement of the frequency of somatic signs. These signs are reliably seen in rats, but are less reliably observed in mice (Malin et al. 1992, 1994; Hildebrand et al. 1997; Epping-Jordan et al. 1998; Isola et al. 1999; Carboni et al. 2000; Semenova et al. 2003). The most prominent signs of the rat withdrawal syndrome include abdominal constrictions (writhes), gasps, ptosis, facial fasciculation, and eyeblinks, while miscellaneous other signs, such as escape attempts, foot licks, genital grooming, shakes, scratches and yawns are rarely observed (Malin et al. 1998; Hildebrand et al. 1997; Watkins et al. 2000b; Cryan et al. 2003). These somatic signs are both centrally and peripherally mediated (Malin et al. 1997; Hildebrand et al. 1997; Watkins et al. 2000a; Carboni et al. 2000). Although the somatic components of nicotine withdrawal are certainly unpleasant, it is hypothesized that avoidance of the affective components of withdrawal plays a more important role in the maintenance of nicotine dependence than the somatic aspects of withdrawal (Watkins et al. 2000b; Kenny & Markou 2001; Hughes 1992). Other rodent models that may be of relevance to disruption of behavioral performance in humans involve nicotine abstinence-induced disruptions of food-maintained learned behaviors in rats (Carroll et al. 1989), increases in the acoustic startle response in rats (Helton et al. 1993), and decreased prepulse inhibition in mice (Semenova et al. 2003). A critical measure of the affective and motivational aspects of drug withdrawal is elevations in brain reward thresholds observed after cessation of chronic nicotine

administration (Epping-Jordan et al. 1998; Harrison et al. 2001; Semenova & Markou 2003; Skjei & Markou 2003; Cryan et al. 2003). Elevations in brain reward thresholds are considered to be an operational measure of “diminished interest or pleasure” in rewarding stimuli that is a symptom of nicotine withdrawal (American Psychiatric Association 1994; Covey et al. 1997). Similar reward threshold elevations are seen during withdrawal from all major drugs of abuse, such as cocaine (Markou & Koob 1991, 1992; Markou et al. 1992; Baldo et al. 1999; Kokkinidis et al. 1980; Frank et al. 1992), ethanol (Schultheis et al. 1995), morphine (Schultheis et al. 1994), amphetamine (Lin et al. 1999, 2000; Paterson et al. 2000; Harrison et al. 2001; Semenova & Markou 2003; Kokkinidis et al. 1980) and phencyclidine (Spielewoy & Markou 2003). Interestingly, several dissociations have been observed between the threshold elevations and the somatic signs associated with nicotine withdrawal (Watkins et al. 2000b; Harrison et al. 2001; Semenova & Markou 2003), suggesting that the various aspects of withdrawal are mediated by different substrates.

Although the substrates of nicotine withdrawal have not been investigated as extensively as the substrates of the reinforcing effects of acute nicotine, there are studies suggesting that adaptations in the circuits mediating the acute effects of nicotine occur with the development of nicotine dependence that lead to the withdrawal syndrome. During nicotine withdrawal precipitated by systemic or intra-VTA administration of the nAChR antagonist mecamylamine in nicotine-treated rats, dopamine dialysate levels were decreased in the nucleus accumbens (Fung et al. 1996; Hildebrand et al. 1998; Carboni et al. 2000) and the central nucleus of the amygdala (Panagis et al. 2000). These mecamylamine injections into the VTA also produced dose-dependently most of the somatic signs of nicotine withdrawal (Hildebrand et al. 1999), suggesting the possible involvement of nAChRs transmission in the VTA in the expression of the somatic signs of nicotine withdrawal. Similar decreases in nucleus accumbens dopamine levels are associated also with withdrawal from other drugs of abuse, such as ethanol, morphine, cocaine and amphetamine (e.g., Rossetti et al. 1992; Weiss et al. 1992). By contrast, increases in dialysate dopamine levels were observed in the frontal cortex (Carboni et al., 2000; however, see Hildebrand et al. 1998), similar to those seen during withdrawal from other drugs of abuse (Acquas & Di Chiara 1992; Bassareo et al. 1995). It is noteworthy that the smoking cessation aid bupropion (an atypical antidepressant; trade name Zyban or Wellbutrin) acts, at least in part, by inhibiting neuronal uptake of dopamine and thereby enhancing dopamine transmission (Terry & Katz 1997; Nomikos et al. 1992). Bupropion has been shown to reverse both the threshold elevations and the somatic signs associated with nicotine withdrawal (Cryan et al. 2003; Malin 2001), while its effects on nicotine self-administration are inconsistent (Shoaib et al. 2002; Glick et al. 2002; Bruijnzeel & Markou 2003).

Considering that these decreases in nucleus accumbens dialysate levels are induced by administration of nAChR antagonists into the VTA, it is suggested that a reduction in endogenous cholinergic tone leads to this effect. The types of nAChRs involved probably include the  $\alpha 4$ -containing high-affinity nAChRs (Harvey & Luetje 1996; Damaj et al. 1995; Harvey et al. 1996; Wonnacott 1997; Epping-Jordan et al. 1998) but not the  $\alpha 7$  nAChRs (Markou & Paterson 2001). Administration of the  $\alpha 7$  nAChR antagonist methyllycaconitine did not precipitate either somatic signs or threshold elevations in nicotine-dependent rats suggesting that these receptors may not be involved in the development of nicotine dependence, despite the fact that  $\alpha 7$  receptors undergo rapid desensitization in the presence of concentrations of nicotine achieved in the brains of smokers (Alkondon et al. 2000; Pidoplichko et al. 1997).

Serotonin (5-HT), and the 5-HT<sub>1A</sub> receptor in particular, also appear to play a role in nicotine withdrawal (Benwell et al. 1990; Kenny et al. 2001). Systemic administration of 5-HT<sub>1A</sub> receptor agonists such as 8-OH-DPAT exacerbated the increased startle response observed during nicotine withdrawal, whereas 5-HT<sub>1A</sub> receptor antagonists, such as WAY-100635, alleviated this enhanced response (Rasmussen et al. 1997; 2000). Further, the responsiveness to 8-OH-DPAT of dorsal raphe nucleus neurons was increased during nicotine withdrawal (Rasmussen & Czachura 1997). Thus, nicotine withdrawal may increase the inhibitory influence of somatodendritic 5-HT<sub>1A</sub> autoreceptors located within the raphe nuclei and thereby decrease 5-HT release into forebrain and limbic brain sites (e.g. Benwell & Balfour 1979, 1982a; Ridley & Balfour 1997). This conclusion is supported by the observation that a serotonergic antidepressant treatment involving the co-administration of the selective serotonin reuptake inhibitor fluoxetine and the 5-HT<sub>1A</sub> receptor antagonist p-MPPI rapidly reversed the elevation in brain-stimulation reward thresholds observed in rats undergoing nicotine withdrawal but did not block the somatic signs of withdrawal (Harrison et al. 2001). These data provide further evidence for a dissociation of the mechanisms mediating affective and somatic aspects of nicotine withdrawal. Consistent with the above, the 5-HT<sub>1A</sub> receptor partial agonist buspirone shows efficacy in smoking cessation trials and may reduce withdrawal severity in abstinent smokers (Hilleman et al. 1992, 1994; West et al. 1991; but see Schneider et al. 1996).

Another transmitter system likely to be involved in nicotine withdrawal is glutamate, considering the role of glutamate in the reinforcing effects of nicotine (see above). Based on the stimulatory effects of glutamate neurotransmission on dopamine release (Mansvelder & McGehee 2000; Schilstrom et al. 1998a), it may be hypothesized that decreased glutamate transmission mediates nicotine withdrawal. Indeed, increased glutamate transmission via antagonism of presynaptic inhibitory Group II metabotropic glutamate receptors (for review, see Cartmell & Schoepp, 2000) reversed the affective aspects of nicotine withdrawal (Kenny et al. 2003). Nevertheless, an agonist at Group II metabotropic glutamate receptors that decreases glutamate

transmission ameliorated the increased startle response seen in rats undergoing nicotine withdrawal (Helton et al. 1997). This result suggests that either decreased or increased glutamate transmission is involved in different aspects of nicotine withdrawal. Further studies are warranted to further investigate the role of glutamate in various aspects of nicotine withdrawal in different brain sites.

In terms of the CRF system and the HPA axis, generally smokers have elevated levels of plasma cortisol (Kirschbaum et al. 1992; Frederick et al. 1998) that do not change significantly within the first several days of quitting smoking (Pickworth et al. 1996; Benowitz et al. 1984; Teneggi et al. 2002) but decline after several weeks of abstinence (Puddey et al. 1984; Frederick et al. 1998). Further, there is rapid tolerance to the effects of nicotine on plasma corticosterone (Benwell & Balfour 1979; Cam & Bassett 1984; Sharp and Bayer 1986). During nicotine withdrawal, both increases and decreases in corticosterone levels have been reported in experimental subjects (Benwell & Balfour 1982b; Rasmussen 1998; Andersson et al. 1989). Further research is warranted to best characterize the effects of nicotine withdrawal in CRF transmission and HPA axis function.

Finally, the opioid system also may be involved in nicotine withdrawal. The opioid receptor agonist morphine reversed withdrawal signs in rats undergoing spontaneous nicotine withdrawal (Malin et al. 1993), while nicotine reduced naloxone-precipitated opiate withdrawal in rats (Zarrindast & Farzin 1996). Further,  $\mu$  opioid receptor knockout mice do not show nicotine place preference or nicotine withdrawal (Berrendero et al. 2002). These results suggest that common neurobiological substrates may mediate nicotine and opiate withdrawal. Accordingly, naloxone and an analog of the endogenous antiopiate, neuropeptide FF, precipitated somatic withdrawal signs after chronic nicotine treatment (Malin et al. 1993, 1996; Carboni et al. 2000; however, see Watkins et al. 2000b) but only with extremely high naloxone doses (Carboni et al. 2000; Malin et al. 1993) compared to those required to precipitate withdrawal in opioid-dependent rats (Gellert & Sparber 1977; Brady & Holtzman 1981; Koob et al. 1989; Higgins & Sellers 1994; Schulteis et al. 1994). Further, naloxone injections did not differentially elevate brain reward thresholds in nicotine- and saline-treated rats (Watkins et al. 2000b). By contrast, low naloxone doses led to conditioned place aversions in nicotine-dependent rats (Ise et al., 2000; Watkins et al. 2000b). Overall, this pattern of results suggests that somatic withdrawal signs and brain reward thresholds are not particularly sensitive to alterations in opioid transmission, while conditioned motivational states may be. Interestingly, both naloxone (Tome et al. 2001) and naltrexone (Almeida et al. 2000) recently have been shown to antagonize nAChRs, suggesting that opioid receptor antagonists may precipitate nicotine withdrawal, at least in part, by directly blocking nAChRs.

### **Chronic Nicotine, Opponent Process, Allostasis and the Development of Dependence**

There are similarities and differences between nicotine and other major drugs of abuse. Although nicotine is an effective reinforcer in animal models, when compared directly to other drugs of abuse such as cocaine, nicotine appears to have less reinforcing efficacy in nondependent animals (Risner & Goldberg, 1983). The paradox, however, is that chronic administration of nicotine produces a robust dependence from the perspective of disruption of brain reward systems. Brain reward thresholds in rats are elevated for periods up to a month with chronic nicotine exposure (Skjei & Markou 2003), probably reflecting a protracted dysphoric state, similarly to the one that leads to relapse in humans. To date there is little or no evidence of an increase in the efficacy of nicotine as a reinforcer in animal models after chronic administration of nicotine sufficient to dramatically alter reward thresholds, but such studies remain to be conducted.

From a motivational perspective, such a shift in reward processing may reflect changes in the dynamics of the reward system associated with opponent and allostatic mechanisms. Solomon and Corbit (1974) postulated that hedonic, affective or emotional states, once initiated, are automatically modulated by the central nervous system with mechanisms that reduce the intensity of hedonic feelings. Termed the opponent-process theory of motivation, Solomon argued that there is affective or hedonic habituation (or tolerance) and affective or hedonic withdrawal (abstinence). More specifically, an *a-process*, which could be either a positive or negative hedonic response, was proposed to 1) occur shortly after presentation of a stimulus, 2) correlate closely with the stimulus intensity, quality and duration of the reinforcer, and 3) show tolerance. In contrast, the *b-process* was proposed 1) to appear after the *a-process* terminated and be slow in onset, 2) to be slow to build up to asymptote and slow to decay, and 3) to get larger with repeated exposure.

From a neurobehavioral perspective it was hypothesized that in brain motivational systems the initial acute effect of an emotional stimulus or drug is opposed or counteracted by homeostatic changes in brain systems. This affect control system was conceptualized as a single negative feedback or an opponent loop that opposes the stimulus-aroused affective state (Solomon & Corbit, 1974; Siegel, 1975; Poulos & Cappell, 1991). In the context of drug dependence, Solomon argued that the first few self-administrations of an opiate drug produce a circumscribed pattern of motivational changes. The onset of the drug effect produces euphoria that is the *a-process*, and this euphoria is followed by a decline in its intensity. Then after the drug wears off, a *b-process* state emerges as an aversive craving state.

More recently, opponent process theory has been expanded into the domains of the neurocircuitry and neurobiology of drug addiction from a physiological perspective (Koob & Le Moal, 2001). An allostatic model of the brain motivational systems was proposed that explains

the persistent changes in motivation that are associated with vulnerability to relapse in addiction. *Allostasis* is defined as the maintenance of stability outside of the normal homeostatic range, where an organism must vary all the parameters of its physiological systems to match them appropriately to chronic demands (e.g., reset the system parameters at a new set point; Sterling & Eyer 1988). Thus, allostasis refers to the integrative adaptive processes maintaining stability through change, a stability that is not within the normal homeostatic range. In this framework, addiction is conceptualized as a continuous dysregulation of brain reward systems that progressively increases resulting in compulsive drug use. Counteradaptive processes such as opponent-process that are part of the normal homeostatic limitation of reward function fail to return within the normal homeostatic range and are hypothesized to form an allostatic state. An allostatic state can be defined as a state of chronic deviation of the regulatory systems from their normal state of operation with establishment of a new set point. This allostatic state that is hypothesized to be reflected in a chronic deviation of the reward set point is fueled not only by dysregulation of reward circuits per se, but also by recruitment of brain and hormonal stress responses.

In terms of nicotine dependence, nicotine, like other drugs of abuse may be hypothesized to drive the brain reward system into an allostatic state. Although there is a less dramatic *a-process* than with other drugs of abuse, there appears to be a protracted *b-process* with nicotine. At least two neurobiological hypotheses can be generated to explain an opponent process/allostatic view of nicotine addiction. At the molecular level, Dani and Heinemann (1996) proposed a desensitization model where nicotine stimulates the mesolimbic dopamine system via activation of nAChRs to produce the reinforcing effects of acute nicotine, but with continued use the inactivation of these receptors by desensitization leads to adaptive tolerance. During abstinence, nicotine levels fall, and the increased number of nAChRs throughout the brain begin to recover to a responsive state that may be dependent on the nAChR subtype. Engaging nAChRs in non-reward-related pathways could contribute to the aversive emotional states associated with nicotine withdrawal. According to this hypothesis, smokers in effect would be ultimately medicating themselves with nicotine to regulate the number of functional nAChRs (Dani & Heinemann, 1996). Recent work shows that such changes may be evident in the  $\alpha$  subunit family, with the  $\alpha 4$ ,  $\alpha 2$  and  $\alpha 7$  subunits showing inactivation with chronic nicotine, but the  $\alpha 3$  subunit (which is similar in structure to the  $\alpha 6$  subunit) showing resistance to desensitization (Olale et al. 1997). Thus, one could speculate that there are two powerful forces for development of nicotine addiction at the receptor level: desensitization of nAChRs in non-reward pathways (that leads to self-medication with nicotine), and resistance to desensitization in reward pathways (positive reinforcement).

An alternative hypothesis at the neurobiological level of analysis is that chronic nicotine exposures "deplete" neurotransmitter systems implicated in the reinforcing effects of nicotine and reward in general. Decreases in dopamine, opioid peptides, serotonin and glutamate all have been observed following chronic nicotine exposure (see above). Brain sites hypothesized to be implicated in these neurochemical dysregulations would include elements of the ventral midbrain such as the VTA and the extended amygdala (central nucleus of the amygdala, bed nucleus of the stria terminalis, and medial nucleus accumbens). However, with other drugs of abuse, a second powerful neurochemical dysregulation of the brain motivational systems has been implicated in the allostatic changes associated with the development of dependence (e.g., recruitment of activity in the hormonal and brain stress systems). Nicotine may be unique in that there is little evidence of a robust activation of the HPA axis response to stress during withdrawal from nicotine, yet there are behavioral signs of anxiety in humans and anxiety-like responses in animal models (Pickworth et al. 1996). How the brain stress systems acting in the brain independent of the HPA axis (such as CRF and neuropeptide Y) respond during nicotine dependence is a challenge for future research.

In summary, nicotine is a somewhat unique drug of abuse in that it produces only a relatively small reinforcing effect acutely, but it certainly has high dependence-inducing efficacy. This dissociation suggests that nicotine may be selectively potentiating some part of the motivational systems associated with opponent processes leading to an allostatic disruption of reward function. Desensitization of brain nAChRs at the molecular level, depletion of brain reward neurotransmitters and disruption of brain stress systems are hypothesized to contribute to such an allostatic state.

## **POLICY IMPLICATIONS FOR TOBACCO CONTROL**

The understanding that tobacco dependence has a pathophysiological basis, which has been increasingly elucidated by advances in understanding the neurobiological mechanisms of nicotine reinforcement and dependence, has implications for tobacco control efforts. Additionally, this understanding strengthens the rationale for comprehensive tobacco programs that emphasize prevention and cessation efforts; just as understanding that the disease of malaria is caused by a mosquito-carried parasite strengthens the importance of comprehensive efforts to prevent exposure, and treat those afflicted to the greatest extent possible.

### **Prevention**

Prevention of tobacco use is critically important because exposure to tobacco is not an innocuous behavior but rather a behavior that triggers a cascade of neurobiological events which can, in turn, affect subsequent behavior. Moreover, each subsequent exposure to tobacco leads to the establishment of tolerance, physiological dependence and a potential strengthening of the



biologically rewarding effects of nicotine. This cascade of events can lead to increased daily self-administration and progression of the dependence process. It is not known if all nicotine-induced changes in brain function, such as 300% increases in nAChR numbers in some brain regions (Perry et al. 1999), and alterations of brain nicotine reinforcement systems (Mansvelder et al. 2002), are fully reversed after nicotine abstinence in all persons. It is plausible that persisting alterations may confer a continuing need for nicotine in some individuals (Henningfield & Slade 1998). Because dependence occurs on a continuum of severity, efforts should be made to divert the usual trajectory of increased tobacco use away from the course to addiction as early as possible to prevent severe addiction from being established. Making every effort to prevent any use, as well as to prevent the development of severe addiction in those who do initiate use, is especially important in developing countries where resources for treating severely addicted tobacco users may be very limited.

Evidence indicates that while all adults are susceptible to the biological effects of tobacco, it also appears plausible that earlier onset of use is associated with a higher risk of developing dependence. Therefore, if the population goal is reduced prevalence of tobacco use, then efforts should be made to discourage initiation of use by persons of all ages (Henningfield et al. 2003). The possibility that substances in tobacco products and tobacco smoke contribute to addiction (Royal College of Physicians of London 2000; Henningfield & Zeller 2003) increases concerns about the risks of tobacco exposure. An additional concern is efforts claiming to reduce the risks of heavy tobacco use by the promotion of products assumed to be less toxic (e.g., Bates et al. 2003), because such promotion carries the risk of undermining prevention and cessation efforts (Henningfield & Fagerstrom 2001; Henningfield et al. 2002, 2003).

## **Cessation**

The major implication of understanding that there is a pathophysiological basis for tobacco addiction also may be considered analogous to the understanding that there is a physiological basis for malaria. Specifically, this understanding implies that although some severely addicted people will be able to achieve lasting cessation without medicine, others may require medicines to enable their nervous and endocrine systems to function satisfactorily in the absence of tobacco-delivered nicotine. It also implies that there are physiological targets through which medications can exert effects that would aid cessation, and that some people may require that the physiological aspects of addiction be treated if they are to sustain cessation.

It is clear that many people can achieve cessation without formal treatment intervention and this is encouraging for developed and developing countries alike. Unfortunately, it is equally clear that many people, particularly heavier tobacco users who are also at greater risk of disease, are unable to sustain cessation without treatment (USDHHS 2000; Royal College of Physicians of

London 2000; Fiore et al. 2000). Because it is not known what percentage of tobacco users require treatment to sustain cessation or which tobacco users benefit the most from specific treatments, a goal of comprehensive tobacco control should be to ensure access to a diversity of effective treatments. An implication for the development of new medications for treating tobacco dependence is that a variety of potential mechanisms have emerged as potential targets for treatment drugs. These include drugs that bind to nAChR subtypes and modulate neurohormonal systems, including those involving dopamine and other catecholamines, serotonin, GABA and glutamate.

It should be emphasized that understanding of tobacco use as a behavior involving a neurobiological driving force does not lessen the importance of non-biological determinants of tobacco addiction. Clearly, factors including social acceptability, knowledge of harmfulness, restrictions of tobacco use, cost, and accessibility affect the risk of developing and sustaining dependence, as well as the probability of cessation (USDHHS 2000; Royal College of Physicians of London 2000; Fiore et al. 2000). The above is consistent with the recommendations of the U.S. Centers for Disease Control and Prevention advocating that comprehensive tobacco control strategies address all such factors and provide access to treatment.

### **Pharmacotherapy – focus on smokeless tobacco**

The treatment of cigarette dependence has been well studied, and a number of behavioral and pharmacologic approaches have been developed and shown to be efficacious. In contrast, much less research has been conducted on the treatment of smokeless tobacco dependence, and the approaches generally have been modeled after smoking cessation treatments. Following is a brief review of the behavioral and pharmacologic approaches that have been studied for smokeless tobacco use.

Hatsukami and Severson (1999) have reviewed the behavioral approaches that have been studied. As with smoking cessation, studies generally have shown that behavioral interventions have improved smokeless tobacco cessation rates compared to control groups with no intervention. Further, there is a dose-response relationship in that more intensive therapy tends to improve rates compared to less intensive interventions. In addition, brief advice from a health care professional can improve smokeless cessation rates, particularly in the dental setting.

There have been few studies of the efficacy of nicotine replacement medications (NRT) for the treatment of smokeless tobacco dependence. Nicotine gum would be the obvious choice of NRT formulations because it would allow the smokeless user to receive the oral stimulation and nicotine normally obtained by tobacco. However, three studies of 2 mg nicotine gum showed minimal efficacy in long-term cessation rates (Sinusas & Coroso 1993; Boyle et al. 1993; Hatsukami et al. 1996, 2000). The general lack of efficacy of the gum may be dose-related, as

there has been little published work on the efficacy of 4 mg nicotine gum on smokeless tobacco use. Moist snuff delivers at least as much nicotine during use as cigarette smoking with peak plasma nicotine levels during Copenhagen use reaching over 20 ng/ml (Fant et al. 1999). Whereas 4 mg nicotine gum would not produce the nicotine levels associated with smokeless tobacco use, it would lead to levels higher than those achieved during 2 mg use. Given that smoking cessation with nicotine gum is dose-related, one could predict that the same might be true for smokeless tobacco cessation.

The transdermal nicotine patch in theory could treat smokeless tobacco dependence by breaking the habit of oral dosing while reducing the nicotine withdrawal symptoms associated with abrupt withdrawal from smokeless tobacco use. The data thus far, however, are equivocal. One study compared transdermal nicotine and mint snuff in a 2x2 factorial design and found that the nicotine patch was effective in increasing short-term abstinence over the placebo patch, and in reducing craving and withdrawal signs and symptoms from spit tobacco. There have been no studies of other NRT formulations (e.g., nasal spray, inhaler, and lozenge). The use of lozenge would be predicted to mimic the effects of nicotine gum because the delivery of nicotine would be comparable in terms of both speed and magnitude. Hatsukami & Severson suggest that nicotine nasal spray would not be an ideal treatment option for smokeless users because the delivery of nicotine from spray would be very rapid in contrast to the slower delivery of nicotine from smokeless tobacco. The nicotine inhaler involves puffing a device that has the general appearance of a cigarette and delivers very low doses of nicotine, and thus would appear less likely to benefit smokeless users.

Two recent studies have shown bupropion to be effective for the treatment of smokeless tobacco dependence. Results from a placebo-controlled trial of 68 smokeless tobacco users indicated that at the end of 12 weeks of therapy, the point-prevalence tobacco abstinence rate was 44% in the bupropion group and 26% in the placebo group ( $p < 0.07$ ); however, the effect was not present at 24 weeks (Dale et al. 2002). Bupropion also was associated with a reduction in nicotine withdrawal symptoms and weight gain. In the second study ( $N=70$ ), bupropion produced significantly higher quit rates for smokeless tobacco cessation at the end of treatment (7 weeks) than placebo ( $p < 0.05$ ) with an odds ratio of 2.73 (Glover et al. 2002). These results suggest that bupropion may be effective in increasing rates of abstinence among smokeless users; however, larger studies are clearly needed to confirm these findings.

In conclusion, significant advances in our understanding of the neurobiology of nicotine reinforcement, dependence and withdrawal provides us with excellent starting points on which to base our efforts to develop effective pharmacological and behavioral interventions to both prevent and treat dependence of smoked and smokeless tobacco products. Further research is needed to continue the investigations of the neurosubstrates mediating not only nicotine addiction but also

the effects of other tobacco ingredients that contribute also to the tobacco use habit, and the synergistic effects of these tobacco ingredients with nicotine that promote this deadly habit.

## **ACKNOWLEDGEMENTS and DISCLOSURES**

Preparation of this manuscript was supported, in part, by NIDA grant DA11946 (AM) Tobacco-Related Disease Research Program grant from the State of California (12RT-0231 to AM), a Novartis Research grant (AM.), NIDA grant DA04398 (GFK), and a Robert Wood Johnson Foundation Innovators Combating Substance Abuse Award (JEH). JEH provides consulting services regarding treatments for tobacco dependence to GlaxoSmithKline Consumer Health Care through Pinney Associates, has a financial interest in a nicotine replacement product under development, and serves as an expert witness in litigation against the tobacco industry by the U.S. Department of Justice and other plaintiffs. We would like to thank Dr. Paul Kenny with assistance with the nAChR section and Mr. Brook Henry with assistance with the CRF section. We greatly appreciate the editorial assistance of Dr. Reginald Fant, whose efforts were supported by the Innovators award to JEH, and Mr. Mike Arends from The Scripps Research Institute. This is publication 15583-NP from The Scripps Research Institute.

## **REFERENCES**

Acquas E, Di Chiara G (1992) Depression of mesolimbic dopamine transmission and sensitization to morphine during opiate abstinence. *J Neurochem* 58:1620-1625.

Addolorato G, Caputo F, Capristo E, Colombo G, Gessa GL, Gasbarrini G (2000) Ability of baclofen in reducing alcohol craving and intake: II. Preliminary clinical evidence. *Alcohol Clin Exp Res* 24:67-71.

Addolorato G, Caputo F, Capristo E, Domenicali M, Bernardi M, Janiri L, Agabio R, Colombo G, Gessa GL, Gasbarrini G (2002a) Baclofen efficacy in reducing alcohol craving and intake: a preliminary double-blind randomized controlled study. *Alcohol* 37:504-508.

Addolorato G, Caputo F, Capristo E, Janiri L, Bernardi M, Agabio R, Colombo G, Gessa GL, Gasbarrini G (2002b) Rapid suppression of alcohol withdrawal syndrome by baclofen. *Am J Med* 112:226-229.

Albuquerque EX, Alkondon M, Pereira EF, Castro NG, Schrattenholz A, Barbosa CT, Bonfante-Cabarcas R, Aracava Y, Eisenberg HM, Maelicke A (1997) Properties of neuronal nicotinic acetylcholine receptors: Pharmacological characterization and modulation of synaptic function. *J Pharmacol Exp Ther* 280:1117-1136.

Alkondon M, Pereira EF, Almeida LE, Randall WR, Albuquerque EX (2000) Nicotine at concentrations found in cigarette smokers activates and desensitizes nicotinic acetylcholine receptors in CA1 interneurons of rat hippocampus. *Neuropharmacology* 39:2726-2739.

Alkondon M, Pereira EF, Barbosa CT, Albuquerque EX (1997) Neuronal nicotinic acetylcholine receptor activation modulates g-aminobutyric acid release from CA1 neurons of rat hippocampal slices. *J Pharmacol Exp Ther* 283:1396-1411.

Almeida LE, Pereira EF, Alkondon M, Fawcett WP, Randall WR, Albuquerque EX (2000) The opioid antagonist naltrexone inhibits activity and alters expression of  $\alpha 7$  and  $\alpha 4\beta 2$  nicotinic receptors in hippocampal neurons: implications for smoking cessation programs, *Neuropharmacology* 39:2740-2755.

American Psychiatric Association (1994) Diagnostic and Statistical Manual of Mental Disorders, 4th edition. American Psychiatric Press, Washington DC.

Andersson K, Fuxe K, Eneroth P, Jansson A, Harfstrand A (1989) Effects of withdrawal from chronic exposure to cigarette smoke on hypothalamic and preoptic catecholamine nerve terminal systems and on the secretion of pituitary hormones in the male rat, *Naunyn Schmiedeberg's Arch Pharmacol* 339:387-396.

Andersson K, Siegel R, Fuxe K, Eneroth P (1983) Intravenous injections of nicotine induce very rapid and discrete reductions of hypothalamic catecholamine levels associated with increases of ACTH, vasopressin and prolactin secretion. *Acta Physiol Scand* 118:35-40.

Araujo DM, Lapchak PA, Collier B, Quirion R (1988) Characterization of N-(<sup>3</sup>H)methylcarbamylcholine binding sites and effect of N-methylcarbamylcholine on acetylcholine release in rat brain. *J Neurochem* 51:292-299.

Arneric SP, Sullivan JP, Williams M (1995) Neuronal nicotinic acetylcholine receptors: Novel targets for CNS therapies. In: Bloom FE, Kupfer D (Eds), *Psychopharmacology: The Fourth Generation of Progress*. Raven Press, New York, pp. 95-110.

Baldo BA, Koob GF, Markou A (1999) Role of adenosine A2 receptors in brain stimulation reward under baseline conditions and during cocaine withdrawal in rats. *J Neurosci* 19:11017-11026.

Balfour DJ (1984) Nicotine and the tobacco smoking habit. In: Balfour DJ (Ed), *International Encyclopedia of Pharmacology and Therapeutics*. Pergamon Press, New York, pp. 61-74.

Balfour DJ, Fagerstrom KO (1996) Pharmacology of nicotine and its therapeutic use in smoking cessation and neurodegenerative disorders. *Pharmacol Ther* 72:51-81.

Bancroft A, Levin ED (2000) Ventral hippocampal alpha4beta2 nicotinic receptors and chronic nicotine effects on memory. *Neuropharmacology* 39:2770-2778.

Bardo MT, Bevins RA, Klebaur JE, Crooks PA, Dwoskin LP (1997) (-)-Nornicotine partially substitutes for (+)-amphetamine in a drug discrimination paradigm in rats. *Pharmacol Biochem Behav* 58:1083-1087.

Bardo MT, Green TA, Crooks PA, Dwoskin LP (1999) Nornicotine is self-administered intravenously by rats. *Psychopharmacology* 146:290-296.

Bassareo V, Tanda G, Di Chiara G (1995) Increase of extracellular dopamine in the medial prefrontal cortex during spontaneous and naloxone-precipitated opiate abstinence. *Psychopharmacology* 122:202-205.

Bates C, Fagerstrom KO, Jarvis M, Kunze M, McNeill A, Ramstrom L (2003) European Union Policy on Smokeless Tobacco, February, 2003. [available from: <http://www.ash.org.uk/html/regulation/html/eusmokeless.html>]

Benowitz NL, Kuyt F, Jacob P 3rd (1984) Influence of nicotine on cardiovascular and hormonal effects of cigarette smoking, *Clin Pharmacol Ther* 36:74-81.

Benwell ME, Balfour DJ (1979) Effects of nicotine administration and its withdrawal on plasma corticosterone and brain 5-hydroxyindoles, *Psychopharmacology* 63:7-11.

Benwell ME, Balfour DJ (1982a) The effects of nicotine administration on 5-HT uptake and biosynthesis in rat brain. *Eur J Pharmacol* 84:71-77.

Benwell ME, Balfour DJ (1982b) Effects of chronic nicotine administration on the response and adaptation to stress, *Psychopharmacology* 76:160-162.

Benwell ME, Balfour DJ, Anderson JM (1990) Smoking-associated changes in the serotonergic systems of discrete regions of human brain. *Psychopharmacology* 102:68-72.

Benwell ME, Balfour DJK (1992) The effects of acute and repeated nicotine treatment on nucleus accumbens dopamine and locomotor activity. *Br J Pharmacol* 105:849-856.

Berrendero F, Kieffer BL, Maldonado R (2002) Attenuation of nicotine-induced antinociception, rewarding effects, and dependence in mu-opioid receptor knock-out mice, *J Neurosci* 22:10935-10940.

Boyle R, Severson H, Lichtenstein E, Gordon J (1993) Smokeless tobacco cessation with nicotine reduction: a placebo controlled trial. Paper presented at the 21st Annual Meeting of the American Public Health Association, San Francisco, CA.

Brady LS, Holtzman SG (1981) Locomotor activity in morphine-dependent and post-dependent rats, *Pharmacol Biochem Behav* 14:361-370.

Browne CL (1990) *The Design of Cigarettes*, 3rd edition. Filter Products Division, Hoechst Celanese Corporation, Charlotte NC.

Bruijnzeel AW, Markou A (2003) Effects of bupropion on responding for nicotine and food under fixed-ratio and progressive ratio schedules of reinforcement in rats, submitted.

Buisson B, Bertrand D (2001) Chronic exposure to nicotine upregulates the human  $\alpha 4\beta 2$  nicotinic acetylcholine receptor function. *J Neurosci* 21:1819-1829.

Buisson B, Bertrand D (2002) Nicotine addiction: The possible role of functional upregulation. *Trends Pharmacol Sci* 23:130-136.

Butschsky M, Bailey D, Henningfield JE, Pickworth WB (1995) Smoking without nicotine delivery decreases withdrawal in 12-hour abstinent smokers. *Pharmacol Biochem Behav* 50:91-96.

Caggiula AR, Donny EC, White AR, Chaudhri N, Booth S, Gharib MA, Hoffman A, Perkins KA, Sved AF (2001a) Cue dependency of nicotine self-administration and smoking. *Pharmacol Biochem Behav* 70:515-530.

Caggiula AR, Donny EC, White AR, Chaudhri N, Booth S, Gharib MA, Hoffman A, Perkins KA, Sved AF (2001b) Environmental stimuli promote the acquisition of nicotine self-administration in rats. *Psychopharmacology* 163:230-237.

Cam GR, Bassett JR (1984) Effect of prolonged exposure to nicotine and stress on the pituitary-adrenocortical response: the possibility of cross-adaptation, *Pharmacol Biochem Behav* 20:221-226.



Cam GR, Bassett JR, Cairncross KD (1979) The action of nicotine on the pituitary-adrenal cortical axis, *Arch Int Pharmacodyn Ther* 237:49-66.

Carboni E, Bortone L, Giua C, DiChiara G (2000) Dissociation of physical abstinence signs from changes in extracellular dopamine in the nucleus accumbens and in the prefrontal cortex of nicotine dependent rats. *Drug Alcohol Depend* 58:93-102.

Carroll ME, Lac ST, Asencio M, Keenan RM (1989) Nicotine dependence in rats. *Life Sci* 45:1381-1388.

Cartmell J, Schoepp DD (2000) Regulation of neurotransmitter release by metabotropic glutamate receptors. *J Neurochem* 75:889-907.

Changeux JP, Devillers-Thiery A, Chemouilli P (1984) Acetylcholine receptor: An allosteric protein. *Science* 225:1335-1345.

Cheeta S, Tucci S, File SE (2001) Antagonism of the anxiolytic effect of nicotine in the dorsal raphe nucleus by dihydro-beta-erythroidine. *Pharmacol Biochem Behav* 70:491-496.

Chen D, Dang H, Patrick JW (1998) Contributions of N-linked glycosylation to the expression of a functional alpha7-nicotinic receptor in *Xenopus* oocytes. *J Neurochem* 70:349-357.

Childress AR, Franklin T, Listerud J, Acton P and O'Brien CP (2002) Neuroimaging of cocaine craving states: Cessation, stimulant administration, and drug cue paradigms [abstract]. College on Problems of Drug Dependence, 64th Annual Meeting.

Childress AR, McLellan AT, Ehrman R, O'Brien CP (1988) Classically conditioned responses in opioid and cocaine dependence: A role in relapse? *NIDA Res Monogr* 84:25-43.

Childress AR, McLellan AT, O'Brien CP (1986) Abstinent opiate abusers exhibit conditioned craving, conditioned withdrawal and reductions in both through extinction. *Br J Addict* 81:655-660.

Churchill L, Dilts RP, Kalivas PW (1992) Autoradiographic localization of gamma-aminobutyric acidA receptors within the ventral tegmental area. *Neurochem Res* 17:101-106.

Clarke PB (1992) The fall and rise of neuronal  $\alpha$ -bungarotoxin binding proteins. *Trends Pharmacol Sci* 13:407-413.

Collins AC, Marks MJ (1989) Chronic nicotine exposure and brain nicotinic receptors: Influence of genetic factors. *Prog Brain Res* 79:137-146.

Colquhoun LM, Patrick JW (1997)  $\alpha 3$ ,  $\beta 2$ , and  $\beta 4$  form heterotrimeric neuronal nicotinic acetylcholine receptors in *Xenopus* oocytes. *J Neurochem* 69:2355-2362.

Conroy WG, Berg DK (1995) Neurons can maintain multiple classes of nicotinic acetylcholine receptors distinguished by different subunit compositions. *J Biol Chem* 270:4424-4431.

Conte-Devolx B, Grino M, Nieoullon A, Javoy-Agid F, Castanas E, Guillaume V, Tonon MC, Vaudry H, Oliver C (1985) Corticoliberin, somatocristin and amine contents in normal parkinsonian human hypothalamus, *Neurosci Lett* 56:217-222.

Corrigall WA, Coen KM (1989) Nicotine maintains robust self-administration in rats on a limited-access schedule. *Psychopharmacology* 99:473-478.

Corrigall WA, Coen KM (1991) Selective dopamine antagonists reduce nicotine self-administration. *Psychopharmacology* 104:171-176.

Corrigall WA, Coen KM, Adamson KL (1994) Self-administered nicotine activates the mesolimbic dopamine system through the ventral tegmental area. *Brain Res* 653:278-284.

Corrigall WA, Coen KM, Adamson KL, Chow BL, Zhang J (2000) Response of nicotine self-administration in the rat to manipulations of  $\mu$ -opioid and gamma-aminobutyric acid receptors in the ventral tegmental area. *Psychopharmacology* 149:107-114.

Corrigall WA, Coen KM, Zhang J, Adamson KL (2001) GABA mechanisms in the pedunculopontine tegmental nucleus influence particular aspects of nicotine self-administration selectively in the rat. *Psychopharmacology* 158:190-197.

Corrigall WA, Franklin KB, Coen KM, Clarke PB (1992) The mesolimbic dopaminergic system is implicated in the reinforcing effects of nicotine. *Psychopharmacology* 107:285-289.

Cousins MS, Stamat HM, de Wit H (2001) Effects of a single dose of baclofen on self-reported subjective effects and tobacco smoking. *Nicotine Tob Res* 3:123-129.

Covey LS, Glassman AH, Stetner F (1997) Major depression following smoking cessation. *Am J Psychiatry* 154:263-265.

Covey LS, Glassman AH, Stetner F (1999) Naltrexone effects on short-term and long-term smoking cessation, *J Addict Dis* 18:31-40.

Crooks PA, Dwoskin LP (1997) Contribution of CNS nicotine metabolites to the neuropharmacological effects of nicotine and tobacco smoking. *Biochem Pharmacol* 54:743-753.

Cryan JF, Bruijnzeel AW, Skjei KL, Markou A (2003) The antismoking and antidepressant agent bupropion enhances brain reward function and reverses the affective and somatic aspects of nicotine withdrawal. *Neuropsychopharmacology*, submitted.

Dale LC, Ebbert JO, Schroeder DR, Croghan IT, Rasmussen DF, Trautman JA, Cox LS, Hurt RD (2002) Bupropion for the treatment of nicotine dependence in spit tobacco users: a pilot study. *Nic Tobacco Res* 4:267-274.

Damaj MI, Glassco W, Dukat M, Martin BR (1999) Pharmacological characterization of nicotine-induced seizures in mice. *J Pharmacol Exp Ther* 291:1284-1291.

Damaj MI, Welch SP, Martin BR (1995) In vivo pharmacological effects of dihydro-beta-erythroidine, a nicotinic antagonist, in mice. *Psychopharmacology* 117:67-73.

Damsma G, Day J, Fibiger HC (1989) Lack of tolerance to nicotine-induced dopamine release in the nucleus accumbens. *Eur J Pharmacol* 168:363-368.

Dani JA (2000) Properties underlying the influence of nicotinic receptors on neuronal excitability and epilepsy. *Epilepsia* 41:1063-1065.

Dani JA, Heinemann S (1996) Molecular and cellular aspects of nicotine abuse. *Neuron* 16:905-908.

Dani JA, Radcliffe KA, Pidoplichko VI (2000) Variations in desensitization of nicotinic acetylcholine receptors from hippocampus and midbrain dopamine areas. *Eur J Pharmacol* 393:31-38.

Deneris ES, Connolly J, Rogers SW, Duvoisin R (1991) Pharmacological and functional diversity of neuronal nicotinic acetylcholine receptors. *Trends Pharmacol Sci* 12:34-40.

Dewey SL, Brodie JD, Gerasimov M, Horan B, Gardner EL, Ashby CR (1999) A pharmacologic strategy for the treatment of nicotine addiction. *Synapse* 31:76-86.

Dewey SL, Morgan AE, Ashby CR Jr, Horan B, Kushner SA, Logan J, Volkow ND, Fowler JS, Gardner EL, Brodie JD (1998) A novel strategy for the treatment of cocaine addiction. *Synapse* 30:119-129.

Dewey SL, Smith GS, Logan J, Brodie JD, Yu DW, Ferrieri RA, King PT, MacGregor RR, Martin TP, Wolf AP, Volkow ND, Fowler JS (1992) GABAergic inhibition of endogenous dopamine release measured in vivo with <sup>11</sup>C-raclopride and positron emission tomography. *J Neurosci* 12:3773-3780.

Donny EC, Caggiula AR, Knopf S, Brown C (1995) Nicotine self-administration in rats. *Psychopharmacology* 122:390-394.

Dwoskin LP, Crooks PA, Teng L, Green TA, Bardo MT (1999) Acute and chronic effects of nornicotine on locomotor activity in rats: altered response to nicotine. *Psychopharmacology* 145:442-451.

Elgoyhen AB, Vetter DE, Katz E, Rothlin CV, Heinemann SF, Boulter J (2001) Alpha 10: A determinant of nicotinic cholinergic receptor function in mammalian vestibular and cochlear mechanosensory hair cells. *Proc Natl Acad Sci USA* 98:3501-3506.

Engberg G, Kling-Petersen T, Nissbrandt H (1993) GABA-B-receptor activation alters the firing pattern of dopamine neurons in the rat substantia nigra. *Synapse* 15:229-238.

Epping-Jordan MP, Watkins SS, Koob GF, Markou A (1998) Dramatic decreases in brain reward function during nicotine withdrawal. *Nature* 393:76-79 .

Fallon JH, Moore RY (1978) Catecholamine innervation of the basal forebrain: IV. Topography of the dopamine projection to the basal forebrain and neostriatum. *J Comp Neurol* 180:545-580.

Fant RV, Henningfield JE, Nelson RA, Pickworth WB (1999) Pharmacokinetics and pharmacodynamics of moist snuff in humans. *Tobacco Control* 8:387-392.

Fattore L, Cossu G, Martellotta MC, Fratta W (2002) Baclofen antagonizes intravenous self-administration of nicotine in mice and rats. *Alcohol Alcohol* 37:495-498.

Fiore MC, Bailey WC, Cohen SJ, Dorfman SF, Goldstein MG, Gritz ER, Heyman RB, Jaen CR, Kottke TE, Lando HA, Mecklenburg RE, Dolan-Mullen P, Nett LM, Robinson L, Stitzer ML, Tommasello AC, Villejo L, Wewers ME (2000) Treating Tobacco Use and Dependence: Clinical Practice Guideline, U.S. Department of Health and Human Services. Public Health Service, Rockville, MD. [available online: <http://www.surgeongeneral.gov/tobacco/>]

Flores CM, Davila-Garcia MI, Ulrich YM, Kellar KJ (1997) Differential regulation of neuronal nicotinic receptor binding sites following chronic nicotine administration. *J Neurochem* 69:2216-2219.

Flores CM, Rogers SW, Pabreza LA, Wolfe BB, Kellar KJ (1992) A subtype of nicotinic cholinergic receptor in rat brain is composed of  $\alpha 4$  and  $\beta 2$  subunits and is up-regulated by chronic nicotine treatment. *Mol Pharmacol* 41:31-37.

Frank RA, Manderscheid PZ, Panicker S, Williams HP, Kokoris D (1992) Cocaine euphoria, dysphoria, and tolerance assessed using drug-induced changes in brain stimulation reward. *Pharmacol Biochem Behav* 42:771-779.

Frederick SL, Reus VI, Ginsberg D, Hall SM, Munoz RF, Ellman G (1998) Cortisol and response to dexamethasone as predictors of withdrawal distress and abstinence success in smokers, *Biol Psychiatry* 43:525-530.

Fu Y, Matta SG, Gao W, Brower VG, Sharp BM (2000) Systemic nicotine stimulates dopamine release in nucleus accumbens: Re-evaluation of the role of N-methyl-D-aspartate receptors in the ventral tegmental area. *J Pharmacol Exp Ther* 294:458-465.

Fudala PJ, Teoh KW, Iwamoto ET (1985) Pharmacologic characterization of nicotine-induced conditioned place preference. *Pharmacol Biochem Behav* 22:237-241.

Fung YK, Schmid MJ, Anderson TM, Lau Y-S (1996) Effects of nicotine withdrawal on central dopaminergic systems. *Pharmacol Biochem Behav* 53:635-640.

Gellert VF, Sparber SB (1977) A comparison of the effects of naloxone upon body weight loss and suppression of fixed-ratio operant behavior in morphine-dependent rats, *J Pharmacol Exp Ther* 201:44-54.

Gentry CL, Wilkins LH Jr, Lukas RJ (2003) Effects of prolonged nicotinic ligand exposure on function of heterologously expressed, human  $\alpha 4\beta 2$ - and  $\alpha 4\beta 4$ -nicotinic acetylcholine receptors. *J Pharmacol Exp Ther* 304:206-216.

Gioanni Y, Rougeot C, Clarke PB, Lepouse C, Thierry AM, Vidal C (1999) Nicotinic receptors in the rat prefrontal cortex: Increase in glutamate release and facilitation of mediodorsal thalamo-cortical transmission. *Eur J Neurosci* 11:18-30.

Glassman AH, Helzer JE, Covey LS, Cottler LB, Stetner F, Tipp JE, Johnson J (1990) Smoking, smoking cessation and major depression. *J Am Med Assoc* 264:1546-1549.

Glick SD, Maisonneuve IM, Kitchen BA (2002) Modulation of nicotine self-administration in rats by combination therapy with agents blocking  $\alpha 3\beta 4$  nicotinic receptors. *Eur J Pharmacol* 448:185-191.

Glover ED, Glover PN, Sullivan CR, Cerullo CL, Hobbs G (2002) A comparison of sustained-release bupropion and placebo for smokeless tobacco cessation. *Am J Health Behav* 26:386-393.

Goldberg SR, Henningfield JE (1988) Reinforcing effects of nicotine in humans and experimental animals responding under intermittent schedules of i.v. drug injection. *Pharmacol Biochem Behav* 30:227-234.

Goldberg SR, Spealman RD, Goldberg DM (1981) Persistent behavior at high rates maintained by intravenous self-administration of nicotine. *Science* 214:573-575.

Goldberg SR, Spealman RD, Risner ME, Henningfield JE (1983) Control of behavior by intravenous nicotine injections in laboratory animals. *Pharmacol Biochem Behav* 19:1011-1020.

Gordon J (1999) AR-R17779: A high affinity, subtype-selective full agonist of the alpha-7 nicotinic acetylcholine receptor. Paper presented at the 2<sup>nd</sup> International Symposium on Nicotinic Acetylcholine Receptors, Annapolis MD, May 13-14, 1999.

Grady SR, Meinerz NM, Cao J, Reynolds AM, Picciotto MR, Changeux JP, McIntosh JM, Marks MJ, Collins AC (2001) Nicotinic agonists stimulate acetylcholine release from mouse interpeduncular nucleus: A function mediated by a different nAChR than dopamine release from striatum. *J Neurochem* 76:258-268.

Gray R, Rajan AS, Radcliffe KA, Yakehiro M, Dani JA (1996) Hippocampal synaptic transmission enhanced by low concentrations of nicotine. *Nature* 383:713-716.

Grenhoff J, Aston-Jones G, Svensson TH (1986) Nicotinic effects on the firing pattern of midbrain dopamine neurons. *Acta Physiol Scand* 128:351-358.

Grillner P, Svensson TH (2000) Nicotine-induced excitation of midbrain dopamine neurons in vitro involves ionotropic glutamate receptor activation. *Synapse* 38:1-9.

Grottick AJ, Trube G, Corrigall WA, Huwyler J, Malherbe P, Wyler R, Higgins GA (2000) Evidence that nicotinic  $\alpha 7$  receptors are not involved in the hyperlocomotor and rewarding effects of nicotine. *J Pharmacol Exp Ther* 294:1112-1119.

Harrison AA, Liem TB, Markou A (2001) Fluoxetine combined with a serotonin-1A receptor antagonist reversed reward deficits observed during nicotine and amphetamine withdrawal in rats. *Neuropsychopharmacology* 25:55-71.

Harvey SC, Leutje CW (1996) Determinates of competitive antagonist sensitivity on neuronal nicotinic receptor beta subunits. *J Neurosci* 16:3798-3806.

Harvey SC, Maddox FN, Luetje CW (1996) Multiple determinants of dihydro-beta-erythroidine sensitivity on rat neuronal nicotinic receptor alpha subunits. *J Neurochem* 67:1953-1959.

Hatsukami D, Jensen J, Allen S, Grillo M, Bliss R (1996) Effects of behavioral and pharmacological treatment on smokeless tobacco users. *J Consult Clin Psychol* 64:153-161.

Hatsukami DK, Grillo M, Boyle R, Allen S, Jensen J, Bliss R, Brown S (2000) Treatment of spit tobacco users with transdermal nicotine system and mint snuff. *J Consult Clin Psychol* 68:241-249.

Hatsukami DK, Severson HH (1999) Oral spit tobacco: addiction, prevention and treatment. *Nicotine Tob Res* 1:21-44.

Heimer L, Alheid GF (1991) Piecing together the puzzle of basal forebrain anatomy. In: Napier TC, Kalivas PW, Hanin I (Eds), *The Basal Forebrain: Anatomy to Function*. Plenum Press, New York, pp. 1-42.

Heimer L, Zahm DS, Churchill L, Kalivas PW, Wohltmann C (1991) Specificity in the projection patterns of accumbal core and shell in the rat. *Neuroscience* 41:89-125.

Heishman SJ, Henningfield JE (2000) Tolerance to repeated nicotine administration on performance, subjective, and physiological responses in nonsmokers. *Psychopharmacology* 152:321-333.

Heishman SJ, Taylor RC, Henningfield JE (1994) Nicotine and smoking: A review of effects on human performance. *Exp Clin Psychopharmacol* 2:345-395.

Helton DR, Modlin DL, Tizzano JP, Rasmussen K (1993) Nicotine withdrawal: A behavioral assessment using schedule controlled responding, locomotor activity, and sensorimotor reactivity. *Psychopharmacology* 113:205-210.

Helton DR, Tizzano JP, Monn JA, Schoepp DD, Kallman MJ (1997) LY354740: A metabotropic glutamate receptor agonist which ameliorates symptoms of nicotine withdrawal in rats. *Neuropharmacology* 36:1511-1516.



Henningfield J, Rose C, Giovino G (2002) Brave new world of tobacco disease prevention: promoting dual tobacco-product use? *Am J Prevent Med* 23:226-228.

Henningfield JE, Fagerstrom KO (2001) Swedish Match Company, Swedish snus and public health: a harm reduction experiment in progress? *Tobacco Control* 10:253-257.

Henningfield JE, Keenan RM, Clarke PBS (1996) Nicotine. In: Schuster CR, Kuhar M (Eds), *Pharmacological Aspects of Drug Dependence: Toward an Integrated Neurobehavioral Approach* (series title: *Handbook of Experimental Pharmacology*, vol. 118), Springer, New York, pp. 272-314.

Henningfield JE, Miyasato K, Jasinski DR (1985) Abuse liability and pharmacodynamic characteristics of intravenous and inhaled nicotine. *J Pharmacol Exp Ther* 234:1-12.

Henningfield JE, Slade J (1998) Tobacco-dependence medications: public health and regulatory issues. *Food Drug Law J* 53(Suppl.):75-114.

Henningfield JE, Zeller M (2003) Could science-based regulation make tobacco products less addictive. *Yale J Health Policy, Law Ethics*, in press.

Henningfield JE, Zeller M, Moolchan E (2003) Regulatory strategies to reduce tobacco addiction in youth. *Tobacco Control*, in press.

Higgins GA, Sellers EM (1994) Antagonist-precipitated opioid withdrawal in rats: evidence for dissociations between physical and motivational signs, *Pharmacol Biochem Behav* 48:1-8.

Hildebrand BE, Nomikos GG, Bondjers C, Nisell M, Svensson TH (1997) Behavioral manifestations of the nicotine abstinence syndrome in the rat: Peripheral versus central mechanisms. *Psychopharmacology* 129:348-356.

Hildebrand BE, Nomikos GG, Hertel P, Schilstrom B, Svensson TH (1998) Reduced dopamine output in the nucleus accumbens but not the medial prefrontal cortex in rats displaying a mecamylamine-precipitated nicotine withdrawal syndrome. *Brain Res* 779:214-225.

Hildebrand BE, Panagis G, Svensson TH, Nomikos GG (1999) Behavioral and biochemical manifestations of mecamylamine-precipitated nicotine withdrawal in the rat: Role of nicotinic receptors in the ventral tegmental area. *Neuropsychopharmacology* 21:560-574.

Hilleman DE, Mohiuddin SM, Del Core MG, Sketch MH (1992) Effect of buspirone on withdrawal symptoms associated with smoking cessation. *Arch Intern Med* 152:350-352.

Hilleman DE, Mohiuddin SM, Delcore MG (1994) Comparison of fixed-dose transdermal nicotine, tapered-dose transdermal nicotine, and buspirone in smoking cessation. *J Clin Pharmacol* 34:222-224.

Hillhouse EW, Milton NG (1989) Effect of noradrenaline and gamma-aminobutyric acid on the secretion of corticotrophin-releasing factor-41 and arginine vasopressin from the rat hypothalamus in vitro, *J Endocrinol* 122:719-723.

Holladay MW, Dart MJ, Lynch JK (1997) Neuronal nicotinic acetylcholine receptors as targets for drug discovery. *J Med Chem* 40:4169-4194.

Hughes JR (1992) Tobacco withdrawal in self-quitters. *J Consult Clin Psychol* 60:689-697.

Hughes JR, Gulliver SB, Fenwick JW, Valliere WA, Cruser K, Pepper S, Shea P, Solomon LJ, Flynn BS (1992) Smoking cessation among self-quitters. *Health Psychol* 11:331-334.

Hughes JR, Gust SW, Keenan R, Fenwick JW, Skoog K, Higgins ST (1991a) Long-term use of nicotine vs placebo gum. *Arch Intern Med* 151:1993-1998.

Hughes JR, Gust SW, Skoog K, Keenan RM, Fenwick JW (1991b) Symptoms of tobacco withdrawal: A replication and extension. *Arch Gen Psychiatry* 48:52-59.

Hughes JR, Hatsukami D (1992) The nicotine withdrawal syndrome: A brief review and update. *Int J Smoking Cessation* 1:21-26.

Hughes JR, Higgins ST, Hatsukami D (1990) Effects of abstinence from tobacco: A critical review. In: Kozlowski LT, Annis HM, Cappell HD, Glaser FB, Goodstadt MS, Israel Y, Kalant

H, Sellers EM, Vingilis ER (Eds), Research Advances in Alcohol and Drug Problems, vol. 10. Plenum, New York, pp. 317-398.

Hunt WA, Barnett LW, Branch LG (1971) Relapse rates in addiction programs. *J Clin Psychol* 27:455-456.

Hunt WA, Bospalec DA (1974) An evaluation of current methods of modifying smoking behavior. *J Clin Psychol* 30:431-438.

Huston-Lyons D, Kornetsky C (1992) Effects of nicotine on the threshold for rewarding brain stimulation in rats. *Pharmacol Biochem Behav* 41:755-759.

Imperato A, Mulas A, DiChiara G (1986) Nicotine preferentially stimulates dopamine release in the limbic system of freely moving rats. *Eur J Pharmacol* 132:337-338.

Ise Y, Narita M, Nagase H, Suzuki T (2000) Modulation of opiodergic system on mecamylamine-precipitated nicotine-withdrawal aversion in rats, *Psychopharmacology* 151:49-54.

Isola R, Vogelsberg V, Wemlinger TA, Neff NH, Hadjiconstantinou M (1999) Nicotine abstinence in the mouse. *Brain Res* 850:189-196.

Jacob P 3rd, Yu L, Shulgin AT, Benowitz NL (1999) Minor tobacco alkaloids as biomarkers for tobacco use: Comparison of users of cigarettes, smokeless tobacco, cigars, and pipes. *Am J Public Health* 89:731-736.

Jung MJ, Lippert B, Metcalf BW, Bohlen P, Schechter PJ (1977) Gamma-vinyl GABA (4-amino-hex-5-enoic acid), a new selective inhibitor of GABA-T: Effects on brain GABA metabolism in mice. *J Neurochem* 29:797-802.

Kaiser SA, Soliakov L, Harvey SC, Luetje CW, Wonnacott S (1998) Differential inhibition by  $\alpha$ -conotoxin-MII of the nicotinic stimulation of ( $^3$ H)dopamine release from rat striatal synaptosomes and slices. *J Neurochem* 70:1069-1076.

Kalivas PW (1993) Neurotransmitter regulation of dopamine neurons in the ventral tegmental area. *Brain Res Rev* 18:75-113.

Kalivas PW, Striplin CD, Steketee JD, Klitenick MA, Duffy P (1992) Cellular mechanisms of behavioral sensitization to drugs of abuse. *Ann NY Acad Sci* 654:128-135.

Karras A, Kane JM (1980) Naloxone reduces cigarette smoking, *Life Sci* 27:1541-1555.

Kasckow JW, Regmi A, Sheriff S, Mulchahey J, Geraciotti TD Jr (1999) Regulation of corticotropin-releasing factor messenger RNA by nicotine in an immortalized amygdalar cell line, *Life Sci* 65:2709-2714.

Kellar KJ, Davila-Garcia MI, Xiao Y (1999) Pharmacology of neuronal nicotinic acetylcholine receptors: Effects of acute and chronic nicotine. *Nicotine Tob Res* 1(Suppl. 2):s117-s120.

Kenny PJ, File SE, Neal MJ (2000) Evidence for a complex influence of nicotinic acetylcholine receptors on hippocampal serotonin release. *J Neurochem* 75:2409-2414.

Kenny PJ, File SE, Rattray M (2001) Nicotine regulates 5-HT<sub>1A</sub> receptor gene expression in the cerebral cortex and dorsal hippocampus. *Eur J Neurosci* 13:1267-1271.

Kenny PJ, Gasparini F, Markou A (2003) Group II metabotropic and alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionate (AMPA)/kainate glutamate receptors regulate the deficit in brain reward function associated with nicotine withdrawal in rats, *J Pharmacol Exp Ther* 306:1068-1076.

Kenny PJ, Markou A (2001) Neurobiology of the nicotine withdrawal syndrome. *Pharmacol Biochem Behav* 70:531-549.

Kirschbaum C, Wust S, Strasburger CJ (1992) "Normal" cigarette smoking increases free cortisol in habitual smokers, *Life Sci* 50:435-442.

Klitenick MA, DeWitte P, Kalivas PW (1992) Regulation of somatodendritic dopamine release in the ventral tegmental area by opioids and GABA: An in vivo microdialysis study. *J Neurosci* 12:2623-2632.

Kokkinidis L, Zacharko RM, Predy PA (1980) Post-amphetamine depression of self-stimulation responding from the substantia nigra: reversal by tricyclic antidepressants. *Pharmacol Biochem Behav* 13:379-383.

Koob GF, Le Moal M (2001) Drug addiction, dysregulation of reward, and allostasis. *Neuropsychopharmacology* 24:97-129.

Koob GF, Markou A, Weiss F, Schulteis G (1993) Opponent process and drug dependence: Neurobiological mechanisms. *Seminars Neurosci* 5:351-358.

Koob GF, Wall TL, Bloom FE (1989) Nucleus accumbens as a substrate for the aversive stimulus effects of opiate withdrawal, *Psychopharmacology* 98:530-534.

Krystal JH, Sanacora G, Blumberg H, Anand A, Charney DS, Marek G, Epperson CN, Goddard A, Mason GF (2002) Glutamate and GABA systems as targets for novel antidepressant and mood-stabilizing treatments. *Mol Psychiatry* 7(Suppl. 1):s71-s80.

Langley JN (1905) On the reaction of cells and of nerve-endings to certain poisons, chiefly as regards the reaction of striated muscle to nicotine and to curari. *J Physiol* 33:374-413.

Lena C, Changeux JP (1998) Allosteric nicotinic receptors, human pathologies. *J Physiol* 92:63-74.

Lena C, de Kerchove D'Exaerde A, Cordero-Erausquin M, Le Novère N, del Mar Arroyo-Jimenez M, Changeux JP (1999) Diversity and distribution of nicotinic acetylcholine receptors in the locus ceruleus neurons. *Proc Natl Acad Sci USA* 96:12126-12131.

Levin ED (1997) Chronic haloperidol administration does not block acute nicotine-induced improvements in the radial-arm maze performance in the rat. *Pharmacol Biochem Behav* 58:899-902.

Levin ED (2002) Nicotinic receptor subtypes and cognitive function. *J Neurobiol* 53:633-640.

Levin ED Simon BB (1998) Nicotine acetylcholine involvement in cognitive function in animals. *Psychopharmacology* 138:217-230.

Levin ED, Bettegowda C, Blosser J, Gordon J (1999) AR-R17779, an  $\alpha 7$  nicotinic agonist, improves learning and memory in rats. *Behav Pharmacol* 10:675-680.

Levine DG (1974) Needle freaks: Compulsive self-injection by drug users. *Am J Psychiatry* 131:297-300.

Lin D, Koob GF, Markou A (1999) Differential effects of withdrawal from chronic amphetamine or fluoxetine administration on brain stimulation reward in the rat: Interactions between the two drugs. *Psychopharmacology* 145:283-294.

Lin D, Koob GF, Markou A (2000) Time-dependent alterations in ICSS thresholds associated with repeated amphetamine administrations. *Pharmacol Biochem Behav* 65:407-417.

Lindstrom J, Anand R, Gerzanich V, Peng X, Wang F, Wells G (1996) Structure and function of neuronal nicotinic acetylcholine receptors. *Prog Brain Res* 109:125-137.

Ling W, Shoptaw S, Majewska D (1998) Baclofen as a cocaine anti-craving medication: A preliminary clinical study. *Neuropsychopharmacology* 18:403-404.

Lippert B, Metcalf BW, Jung MJ, Casara P (1977) 4-amino-hex-5-enoic acid, a selective catalytic inhibitor of 4-aminobutyric-acid aminotransferase in mammalian brain. *Eur J Biochem* 74:441-445.

Ludwig AM (1986) Pavlov's "bells" and alcohol craving. *Addict Behav* 11:87-91.

Luo S, Kulak JM, Cartier GE, Jacobsen RB, Yoshikami D, Olivera BM, McIntosh JM (1998) Alpha-conotoxin AuIB selectively blocks  $\alpha 3\beta 4$  nicotinic acetylcholine receptors and nicotine-evoked norepinephrine release. *J Neurosci* 18:8571-8579.

Malin DH (2001) Nicotine dependence: studies with a laboratory model. *Pharmacol Biochem Behav* 70:551-559.

Malin DH, Lake JR, Carter VA, Cunningham JS, Hebert KM, Conrad DL, Wilson OB (1994) The nicotinic antagonist mecamylamine precipitates nicotine abstinence syndrome in the rat. *Psychopharmacology* 115:180-184.

Malin DH, Lake JR, Carter VA, Cunningham JS, Wilson OB (1993) Naloxone precipitates nicotine abstinence syndrome in the rat, *Psychopharmacology* 112:339-342.

Malin DH, Lake JR, Newlin-Maultsby P, Roberts LK, Lanier JG, Carter VA, Cunningham JS, Wilson OB (1992) Rodent model of nicotine abstinence syndrome. *Pharmacol Biochem Behav* 43:779-784.

Malin DH, Lake JR, Schopen CK, Kirk JW, Sailer EE, Lawless BA, Upchurch TP, Shenoi M, Rajan N (1997) Nicotine abstinence syndrome precipitated by central but not peripheral hexamethonium. *Pharmacol Biochem Behav* 58:695-699.

Malin DH, Lake JR, Short PE, Blossman JB, Lawless BA, Schopen CK, Sailer EE, Burgess K, Wilson OB (1996) Nicotine abstinence syndrome precipitated by an analog of neuropeptide FF, *Pharmacol Biochem Behav* 54:581-585.

Malin DH, Lake JR, Upchurch TP, Shenoi M, Rajan N, Schweinle WE (1998) Nicotine abstinence syndrome precipitated by the competitive nicotinic antagonist dihydro- $\beta$ -erythroidine. *Pharmacol Biochem Behav* 60:609-613.

Mansvelder HD, Keath JR, McGehee DS (2002) Synaptic mechanisms underlie nicotine-induced excitability of brain reward areas. *Neuron* 33:905-919.

Mansvelder HD, McGehee DS (2000) Long-term potentiation of excitatory inputs to brain reward areas by nicotine. *Neuron* 27:349-357.

Markou A, Hauger RL, Koob GF (1992) Desmethylinipramine attenuates cocaine withdrawal in rats. *Psychopharmacology* 109:305-314.

Markou A, Koob GF (1991) Postcocaine anhedonia: An animal model of cocaine withdrawal. *Neuropsychopharmacology* 4:17-26.

Markou A, Koob GF (1992) Bromocriptine reverses the elevation in intracranial self-stimulation thresholds observed in a rat model of cocaine withdrawal. *Neuropsychopharmacology* 7:213-224.

Markou A, Kosten TR, Koob GF (1998) Neurobiological similarities in depression and drug dependence: A self-medication hypothesis. *Neuropsychopharmacology* 18:135-174.

Markou A, Paterson NE (2001) The nicotinic antagonist methyllycaconitine has differential effects on nicotine self-administration and nicotine withdrawal in the rat. *Nicotine Tob Res* 3:361-373.

Markou A, Weiss F, Gold LH, Caine SB, Schulteis G, Koob GF (1993) Animal models of drug craving. *Psychopharmacology* 112:163-182.

Marks MJ, Burch JB, Collins AC (1983) Effects of chronic nicotine infusion on tolerance development and nicotinic receptors. *J Pharmacol Exp Ther* 226:817-825.

Marks MJ, Grady SR, Collins AC (1993) Downregulation of nicotinic receptor function after chronic nicotine infusion. *J Pharmacol Exp Ther* 266:1268-1276.

Marks MJ, Pauly JR, Gross SD, Deneris ES, Hermans-Borgmeyer I, Heinemann SF, Collins AC (1992) Nicotine binding and nicotinic receptor subunit RNA after chronic nicotine treatment. *J Neurosci* 12:2765-2784.

Matta SG, Beyer HS, McAllen KM, Sharp BM (1987) Nicotine elevates rat plasma ACTH by a central mechanism. *J Pharmacol Exp Ther* 243:217-226.

Matta SG, Foster CA, Sharp BM (1993) Selective administration of nicotine into catecholaminergic regions of rat brainstem stimulates adrenocorticotropin secretion, *Endocrinology* 133:2935-2942.



Matta SG, Valentine JD, Sharp BM (1997) Nicotinic activation of CRH neurons in extrahypothalamic regions of the rat brain, *Endocrine* 7:245-253.

McGehee DS, Heath MJ, Gelber S, Devay P, Role LW (1995) Nicotine enhancement of fast excitatory synaptic transmission in CNS by presynaptic receptors. *Science* 269:1692-1696.

McGehee DS, Role LW (1995) Physiological diversity of nicotinic acetylcholine receptors expressed by vertebrate neurons. *Annu Rev Physiol* 57:521-546.

McGehee DS, Role LW (1996) Presynaptic ionotropic receptors. *Curr Opin Neurobiol* 6:342-349.

McLellan AT, Childress AR, Ehrman R, O'Brien CP, Pashko S (1986) Extinguishing conditioned responses during opiate dependence treatment: Turning laboratory findings into clinical procedures. *J Subst Abuse Treat* 3:33-40.

Mifsud JC, Hernandez L, Hoebel BG (1989) Nicotine infused into the nucleus accumbens increased synaptic dopamine as measured by in vivo microdialysis. *Brain Res* 478:365-367.

Nemeth-Coslett R, Griffiths RR (1986) Naloxone does not affect cigarette smoking, *Psychopharmacology* 89:261-264.

Newhouse PA, Sunderland T, Narang PK, Mellow AM, Fertig JB, Lawlor BA, Murphy DL (1990) Neuroendocrine, physiologic, and behavioral responses following intravenous nicotine in nonsmoking healthy volunteers and in patients with Alzheimer's disease, *Psychoneuroendocrinology* 15:471-484.

Nisell M, Marcus M, Nomikos GG, Svensson TH (1997) Differential effects of acute and chronic nicotine on dopamine output in the core and shell of the rat nucleus accumbens. *J Neural Transm* 104:1-10.

Nisell M, Nomikos GG, Svensson TH (1994a) Infusion of nicotine in the ventral tegmental area or the nucleus accumbens of the rat differentially affects accumbal dopamine release. *Pharmacol Toxicol* 75:348-352.

Nisell M, Nomikos GG, Svensson TH (1994b) Systemic nicotine-induced dopamine release in the rat nucleus accumbens is regulated by nicotinic receptors in the ventral tegmental area. *Synapse* 16:36-44.

Nisell M, Nomikos GG, Svensson TH (1995) Nicotine dependence, midbrain dopamine systems and psychiatric disorders. *Pharmacol Toxicol* 76:157-162.

Nomikos GG, Damsma G, Wenkstern D, Fibiger HC (1992) Effects of chronic bupropion on interstitial concentrations of dopamine in rat nucleus accumbens and striatum. *Neuropsychopharmacology* 7:7-14.

O'Brien CP, Childress AR, McLellan AT, Ehrman R (1990) Integrating systematic cue exposure with standard treatment in recovering drug dependent patients. *Addict Behav* 15:355-365.

Okuda H, Shioda S, Nakai Y, Nakayama H, Okamoto M, Nakashima T (1993) The presence of corticotropin-releasing factor-like immunoreactive synaptic immunoreactivity in the median eminence of the rat, *Neurosci Lett* 161:183-186.

Olale F, Gerzanich V, Kuryatov A, Wang F, Lindstrom J (1997) Chronic nicotine exposure differentially affects the function of human alpha-3, alpha-4, and alpha-7 neuronal nicotinic receptor subtypes. *J Pharmacol Exp Ther* 283:675-683.

Panagis G, Hildebrand BE, Svensson TH, Nomikos GG (2000) Selective c-fos induction and decreased dopamine release in the central nucleus of amygdala in rats displaying a mecamylamine-precipitated nicotine withdrawal syndrome. *Synapse* 35:15-25.

Parrott AC (1993) Cigarette smoking: Effects upon self-rated stress and arousal over the day. *Addict Behav* 18:389-395.

Paterson D, Nordberg A (2000) Neuronal nicotinic receptors in the human brain. *Prog Neurobiol* 61:75-111.

Paterson NE, Froestl W, Markou A (2003) The GABA<sub>B</sub> receptor agonists baclofen and CGP44532 decrease nicotine self-administration in the rat. *Neuropharmacology*, under review.

Paterson NE, Markou A (2002) Increased GABA neurotransmission via administration of gamma-vinyl GABA decreased nicotine self-administration in the rat. *Synapse* 44:252-253.

Paterson NE, Myers C, Markou A (2000) Effects of repeated withdrawal from continuous amphetamine administration on brain reward function in rats. *Psychopharmacology* 152:440-446.

Perkins KA, Gerlach D, Vender J, Grobe J, Meeker J, Hutchison S (2001) Sex differences in the subjective and reinforcing effects of visual and olfactory cigarette smoke stimuli. *Nicotine Tob Res* 3:141-150.

Perry DC, Davila-Garcia MI, Stockmeier CA, Kellar KJ (1999) Increased nicotinic receptors in brains from smokers: membrane binding and autoradiography studies. *J Pharmacol Exp Ther* 289:1545-1552.

Picciotto MR, Corrigall WA (2002) Neuronal systems underlying behaviors related to nicotine addiction: Neural circuits and molecular genetics. *J Neurosci* 22:3338-3341.

Picciotto MR, Zoli M, Lena C, Bessis A, Lallemand Y, LeNovere N, Vincent P, Pich EM, Brulet P, Changeux JP (1995) Abnormal avoidance learning in mice lacking functional high-affinity nicotine receptor in the brain. *Nature* 374:65-67.

Picciotto MR, Zoli M, Rimondini R, Lena C, Marubio LM, Merlo-Pich E, Fuxe K, Changeux JP (1998) Acetylcholine receptors containing the beta-2 subunit are involved in the reinforcing properties of nicotine. *Nature* 391:173-177.

Pickworth WB, Baumann MH, Fant RV, Rothman RB, Henningfield JE (1996) Endocrine responses during acute nicotine withdrawal, *Pharmacol Biochem Behav* 55:433-437.

Pickworth WB, Heishman SJ, Henningfield JE (1995) Relationships between EEG and performance during nicotine withdrawal and administration. In: Domino EF (Ed), *Brain Imaging of Nicotine and Tobacco Smoking*, NPP Books, Ann Arbor MI, pp. 1-13.

Pidoplichko VI, DeBiasi M, Williams JT, Dani JA (1997) Nicotine activates and desensitizes midbrain dopamine neurons. *Nature* 390:401-404.

Pomerleau CS, Pomerleau OF (1992) Euphoriant effects of nicotine in smokers. *Psychopharmacology* 108:460-465.

Pontieri FE, Tanda G, Orzi F, DiChiara G (1996) Effects of nicotine on the nucleus accumbens and similarity to those of addictive drugs. *Nature* 382:255-257.

Poulos CX, Cappell H (1991) Homeostatic theory of drug tolerance: A general model of physiological adaptation. *Psychol Rev* 98:390-408.

Puddey IB, Vandongen R, Beilin LJ, English D (1984) Haemodynamic and neuroendocrine consequences of stopping smoking: a controlled study, *Clin Exp Pharmacol Physiol* 11:423-426.

Quick MW, Lester RA (2002) Desensitization of neuronal nicotinic receptors. *J Neurobiol* 53:457-478.

Radcliffe KA, Dani JA (1998) Nicotinic stimulation produces multiple forms of increased glutamatergic synaptic transmission. *J Neurosci* 18:7075-7083.

Rasmussen DD (1998) Effects of chronic nicotine treatment and withdrawal on hypothalamic proopiomelanocortin gene expression and neuroendocrine regulation, *Psychoneuroendocrinology* 23:245-259.

Rasmussen K, Calligaro DO, Czachura JF, Dreshfield-Ahmad LJ, Evans DC, Hemrick-Luecke SK, Kallman MJ, Kendrick WT, Leander JD, Nelson DL, Overshiner CD, Wainscott DB, Wolff MC, Wong DT, Branchek TA, Zgombick JM, Xu YC (2000) The novel 5-hydroxytryptamine<sub>1A</sub> antagonist LY426965: Effects on nicotine withdrawal and interactions with fluoxetine. *J Pharmacol Exp Ther* 294:688-700.

Rasmussen K, Czachura JF (1997) Nicotine withdrawal leads to increased sensitivity of serotonergic neurons to the 5-HT<sub>1A</sub> agonist 8-OH-DPAT. *Psychopharmacology* 133:343-346.

Rasmussen K, Kallman MJ, Helton DR (1997) Serotonin-1A antagonists attenuate the effects of nicotine withdrawal on the auditory startle response. *Synapse* 27:145-152.

Reid MS, Fox L, Ho LB, Berger SP (2000) Nicotine stimulation of extracellular glutamate levels in the nucleus accumbens: Neuropharmacological characterization. *Synapse* 35:129-136.

Ridley DL, Balfour DJK (1997) The influence of nicotine on 5-HT overflow in the dorsal hippocampus of the rat. *Br J Pharmacol* 112(Suppl.):301P.

Risner ME, Goldberg SR (1983) A comparison of nicotine and cocaine self-administration in the dog: Fixed-ratio and progressive-ratio schedules of intravenous drug infusion. *J Pharmacol Exp Ther* 224:319-326.

Role LW, Berg DK (1996) Nicotinic receptors in the development and modulation of CNS synapses. *Neuron* 16:1077-1085.

Rose JE, Behm FM, Levin ED (1993) Role of nicotine dose and sensory cues in the regulation of smoke intake. *Pharmacol Biochem Behav* 44:891-900.

Rossetti ZL, Hmaidan Y, Gessa GL (1992) Marked inhibition of mesolimbic dopamine release: A common feature of ethanol, morphine, cocaine, and amphetamine abstinence in rats. *Eur J Pharmacol* 221:227-234.

Rowell PP, Wonnacott S (1990) Evidence for functional activity of upregulated nicotine binding sites in rat striatal synaptosomes. *J Neurochem* 55:2105-2110.

Royal College of Physicians of London (2000) Nicotine Addiction in Britain: A Report of the Tobacco Advisory Group of the Royal College of Physicians. Royal College of Physicians of London, London.

Royal Society of Canada (1989) Tobacco, Nicotine and Addiction: A Committee Report. Royal Society of Canada, Ottawa.

Sargent PB (1993) The diversity of neuronal nicotinic acetylcholine receptors. *Annu Rev Neurosci* 16:403-443.

Schilstrom B, Nomikos GG, Nisell M, Hertel P, Svensson TH (1998a) N-methyl-D-aspartate receptor antagonism in the ventral tegmental area diminishes the systemic nicotine-induced dopamine release in the nucleus accumbens. *Neuroscience* 82:781-789.

Schilstrom B, Svensson HM, Svensson TH, Nomikos GG (1998b) Nicotine and food induced dopamine release in the nucleus accumbens of the rat: Putative role of  $\alpha 7$  nicotinic receptors in the ventral tegmental area. *Neuroscience* 85:1005-1009.

Schneider NG, Olmstead RE, Steinberg C, Sloan K, Daims RM, Brown HV (1996) Efficacy of buspirone in smoking cessation: A placebo-controlled trial. *Clin Pharmacol Ther* 60:568-575.

Schultheis G, Markou A, Cole M, Koob GF (1995) Decreased brain reward produced by ethanol withdrawal. *Proc Natl Acad Sci USA* 92:5880-5884.

Schultheis G, Markou A, Gold LH, Stinus L, Koob GF (1994) Relative sensitivity to naloxone of multiple indices of opiate withdrawal: A quantitative dose-response analysis. *J Pharmacol Exp Ther* 271:1391-1398.

Schulz DW, Loring RH, Aizenman E, Zigmond RE (1991) Autoradiographic localization of putative nicotinic receptors in the rat brain using  $^{125}\text{I}$ -neuronal bungarotoxin. *J Neurosci* 11:287-297.

Schwartz RD, Kellar KJ (1983) Nicotinic cholinergic receptor binding sites in the brain: Regulation in vivo. *Science* 220:214-216.

Semenova S, Beshpalov A, Markou A (2003) Decreased prepulse inhibition during nicotine withdrawal in DBA/2J mice is reversed by nicotine self-administration. *Eur J Pharmacol*, submitted.

Semenova S, Markou A (2003) Clozapine treatment attenuated somatic and affective signs of nicotine and amphetamine withdrawal in subsets of rats that exhibited hyposensitivity to the initial effects of clozapine. *Biol Psychiatry*, in press.

Sershen H, Balla A, Lajtha A, Vizi ES (1997) Characterization of nicotinic receptors involved in the release of noradrenaline from the hippocampus. *Neuroscience* 77:121-130.

Seyler LE Jr, Fertig J, Pomerleau O, Hunt D, Parker K (1984) The effects of smoking on ACTH and cortisol secretion. *Life Sci* 34:57-65.

Shahan TA, Bickel WK, Madden GJ, Badger GJ (1999) Comparing the reinforcing efficacy of nicotine containing and de-nicotinized cigarettes: A behavioral economic analysis. *Psychopharmacology* 147:210-216.

Sharp BM, Beyer HS (1986) Rapid desensitization of the acute stimulatory effects of nicotine on rat plasma adrenocorticotropin and prolactin. *J Pharmacol Exp Ther* 238:486-491.

Sharples CG, Kaiser S, Soliakov L, Marks MJ, Collins AC, Washburn M, Wright E, Spencer JA, Gallagher T, Whiteaker P, Wonnacott S (2000) UB-165: A novel nicotinic agonist with subtype selectivity implicates the  $\alpha 4\beta 2$  subtype in the modulation of dopamine release from rat striatal synaptosomes. *J Neurosci* 20:2783-2791.

Sheffield EB, Quick MW, Lester RA (2000) Nicotinic acetylcholine receptor subunit mRNA expression and channel function in medial habenula neurons. *Neuropharmacology* 39:2591-2603.

Shiffman S (1989) Trans-situational consistency in smoking relapse. *Health Psychol* 8:471-481.

Shiffman S, Mason KM, Henningfield JE (1998) Tobacco dependence treatments: Review and prospectus. *Annu Rev Public Health* 19:335-358.

Shiffman SM, Jarvik ME (1976) Smoking withdrawal symptoms in two weeks of abstinence. *Psychopharmacology* 50:35-39.

Shoaib M, Benwell ME, Akbar MT, Stolerman IP, Balfour DJ (1994) Behavioural and neurochemical adaptations to nicotine in rats: influence of NMDA antagonists. *Br J Pharmacol* 111:1073-1080.

Shoaib M, Schindler CW, Goldberg SR (1997) Nicotine self-administration in rats: Strain and nicotine pre-exposure effects on acquisition. *Psychopharmacology* 129:35-43.

Shoaib M, Sidhpura N, Shafait S (2002) Investigating the actions of bupropion on dependence-related effects of nicotine in rats. *Psychopharmacology*, in press (<http://link.springer-ny.com/link/service.../01277/paper/s00213-002-1277-xch110.html>).

Shoaib M, Stolerman IP (1992) MK801 attenuates behavioural adaptation to chronic nicotine administration in rats. *Br J Pharmacol* 105:514-515.

Shoaib M, Swanner LS, Beyer CE, Goldberg SR, Schindler CW (1998) The GABAB agonist baclofen modifies cocaine self-administration in rats. *Behav Pharmacol* 9:195-206.

Siegel S (1975) Evidence from rats that morphine tolerance is a learned response. *J Comp Physiol Psychol* 89:498-506.

Sinusas K, Coroso JG (1993) Smokeless tobacco cessation: report of a preliminary trial using nicotine chewing gum. *J Family Practice* 37:264-267.

Skjei KL, Markou A (2003) Effects of repeated withdrawal episodes, nicotine dose, and duration of nicotine exposure on the severity and duration of nicotine withdrawal in rats. *Psychopharmacology*, in press.

Solomon RL, Corbit JD (1974) An opponent-process theory of motivation: 1. Temporal dynamics of affect. *Psychol Rev* 81:119-145.

Soria R, Stapleton JM, Gilson SF, Sampson-Cone A, Henningfield JE, London ED (1996) Subjective and cardiovascular effects of intravenous nicotine in smokers and non-smokers. *Psychopharmacology* 128:221-226.

Spielewoy C, Markou A (2003) Withdrawal from chronic treatment with phencyclidine induces long-lasting depression in brain reward function, *Neuropsychopharmacology*, in press.

Stalke J, Hader O, Bahr V, Hensen J, Scherer G, Oelkers W (1992) The role of vasopressin in the nicotine-induced stimulation of ACTH and cortisol in men. *Clin Investig* 70:218-223.



Sterling P, Eyer J (1988) Allostasis: A new paradigm to explain arousal pathology. In: Fisher S, Reason J (Eds), *Handbook of Life Stress, Cognition and Health*, John Wiley, Chichester, 1988, pp. 629-649.

Stewart J, de Wit H, Eikelboom R (1984) Role of unconditioned and conditioned drug effects in the self-administration of opiates and stimulants. *Psychol Rev* 91:251-268.

Stolerman IP (1991) Behavioral pharmacology of nicotine: Multiple mechanisms. *Br J Addict* 86:533-536.

Stolerman IP, Jarvis MJ (1995) The scientific case that nicotine is addictive. *Psychopharmacology* 117:2-10.

Sugita S, Johnson SW, North RA (1992) Synaptic inputs to GABAA and GABAB receptors originate from discrete afferent neurons. *Neurosci Lett* 134:207-211.

Sutherland G, Stapleton JA, Russell MA, Feyerabend C (1995) Naltrexone, smoking behaviour and cigarette withdrawal, *Psychopharmacology* 120:418-425.

Swedberg MDB, Henningfield JE, Goldberg SR (1990) Nicotine dependency: Animal studies. In Wonnacott S, Russell MAH, Stolerman IP (Eds), *Nicotine Psychopharmacology: Molecular, Cellular, and Behavioural Aspects*, Oxford University Press, New York, pp. 38-76.

Teneggi V, Tiffany ST, Squassante L, Milleri S, Ziviani L, Bye A (2002) Smokers deprived of cigarettes for 72 h: effect of nicotine patches on craving and withdrawal. *Psychopharmacology* 164:177-187.

Teng L, Crooks PA, Buxton ST, Dwoskin LP (1997) Nicotinic-receptor mediation of S(-)-nornicotine-evoked-3H-overflow from rat striatal slices preloaded with [3H]dopamine. *J Pharmacol Exp Ther* 283:778-787.

Terry P, Katz JL (1997) Dopaminergic mediation of the discriminative stimulus effects of bupropion in rats. *Psychopharmacology* 134:201-212.

Toide K, Arima T (1989) Effects of cholinergic drugs on extracellular levels of acetylcholine and choline in rat cortex, hippocampus and striatum studied by brain dialysis. *Eur J Pharmacol* 173:133-141.

Tome AR, Izaguirre V, Rosario LM, Cena V, Gonzalez-Garcia C (2001) Naloxone inhibits nicotine-induced receptor current and catecholamine secretion in bovine chromaffin cells, *Brain Res* 903:62-65.

Toth E, Sershen H, Hashim A, Vizi ES, Lajtha A (1992) Effect of nicotine on extracellular levels of neurotransmitters assessed by microdialysis in various brain regions: Role of glutamic acid. *Neurochem Res* 17:265-271.

US Department of Health and Human Services (1988) The Health Consequences of Smoking: Nicotine Addiction. A Report of the Surgeon General [DHHS publication no. (CDC) 88-8406]. US Government Printing Office, Washington DC.

US Department of Health and Human Services (2000) Reducing Tobacco Use: A Report of the Surgeon General. US Government Printing Office, Washington.

Valentine JD, Matta SG, Sharp BM (1996) Nicotine-induced c-fos expression in the hypothalamic paraventricular nucleus is dependent on brainstem effects: correlations with c-fos in catecholaminergic and noncatecholaminergic neurons in the nucleus tractus solitarius. *Endocrinology* 137:622-630.

Vidal C (1996) Nicotinic receptors in the brain: molecular biology, function, and therapeutics. *Mol Chem Neuropathol* 28:3-11.

Vidal C, Changeux JP (1993) Nicotinic and muscarinic modulations of excitatory synaptic transmission in the rat prefrontal cortex in vitro. *Neuroscience* 56:23-32.

Walaas I, Fonnum F (1979) The distribution and origin of glutamate decarboxylase and choline acetyltransferase in ventral pallidum and other basal forebrain regions. *Brain Res* 177:325-336.

Walaas I, Fonnum F (1980) Biochemical evidence for gamma-aminobutyrate containing fibers from the nucleus accumbens to the substantia nigra and ventral tegmental area in the rat. *Neuroscience* 5:63-72.

Watkins SS, Epping-Jordan MP, Koob GF, Markou A (1999) Blockade of nicotine self-administration with nicotinic antagonists in rats. *Pharmacol Biochem Behav* 62:743-751.

Watkins SS, Koob GF, Markou A (2000a) Neural mechanisms underlying nicotine addiction: acute positive reinforcement and withdrawal. *Nicotine Tob Res* 2:19-37.

Watkins SS, Stinus L, Koob GF, Markou A (2000b) Reward and somatic changes during precipitated nicotine withdrawal in rats: Centrally and peripherally mediated effects. *J Pharmacol Exp Ther* 292:1053-1064.

Weiss F, Markou A, Lorang MT, Koob GF (1992) Basal extracellular dopamine levels in the nucleus accumbens are decreased during cocaine withdrawal after unlimited-access self-administration. *Brain Res* 593:314-318.

West R, Hajek P, McNeill A (1991) Effect of buspirone on cigarette withdrawal symptoms and short-term abstinence rates in a smokers clinic. *Psychopharmacology* 104:91-96.

West RJ, Jarvis MJ, Russell MA, Carruthers ME, Feyerabend C (1984) Effect of nicotine replacement on the cigarette withdrawal syndrome. *Br J Addict* 79:215-219.

Wewers ME, Dhatt R, Tejwani GA (1998) Naltrexone administration affects ad libitum smoking behavior, *Psychopharmacology* 140:185-190.

Wikler A (1973) Dynamics of drug dependence: Implications of a conditioning theory for research and treatment, *Arch Gen Psychiatry* 28:611-616.

Wilkie GI, Hutson P, Sullivan JP, Wonnacott S (1996) Pharmacological characterization of a nicotinic autoreceptor in rat hippocampal synaptosomes. *Neurochem Res* 21:1141-1148.

Wilkins JN, Carlson HE, Van Vunakis H, Hill MA, Gritz E, Jarvik ME (1982) Nicotine from cigarette smoking increases circulating levels of cortisol, growth hormone, and prolactin in male chronic smokers, *Psychopharmacology* 78:305-308.

Williams M, Robinson JL (1984) Binding of the nicotinic cholinergic antagonist, dihydro-beta-erythroidine, to rat brain tissue. *J Neurosci* 4:2906-2911.

Wonnacott S (1990) The paradox of nicotinic acetylcholine receptor upregulation by nicotine. *Trends Pharmacol Sci* 11:216-219.

Wonnacott S (1997) Presynaptic nicotinic ACh receptors. *Trends Neurosci* 20:92-98.

World Health Organization (2001) *Advancing Knowledge on Regulating Tobacco Products*. World Health Organization, Geneva.

Yim CY, Mogenson GJ (1980) Electrophysiological studies of neurons in the ventral tegmental area of Tsai. *Brain Res* 181:301-313.

Zarrindast MR, Farzin D (1996) Nicotine attenuates naloxone-induced jumping behaviour in morphine-dependent mice, *Eur J Pharmacol* 298:1-6.